

MOLECULAR CELL BIOLOGY

T H I R D E D I T I O N


HARVEY LODISH • DAVID BALTIMORE
ARNOLD BERK • S. LAWRENCE ZIPURSKY
PAUL MATSUDAIRA • JAMES DARNELL



Contents in Brief

| | | | |
|---|------------|--|------------|
| ► Part I | | | |
| Laying the Groundwork | 1 | | |
| 1 The Dynamic Cell | 3 | 10 DNA Replication, Repair, and Recombination | 365 |
| 2 Chemical Foundations | 15 | 11 Regulation of Transcription Initiation | 405 |
| 3 Protein Structure and Function | 51 | 12 Transcription Termination, RNA Processing, and Posttranscriptional Control | 485 |
| 4 Nucleic Acids, the Genetic Code, and Protein Synthesis | 101 | 13 Gene Control in Development | 543 |
| 5 Cell Organization, Subcellular Structure, and Cell Division | 141 | ► Part III | |
| 6 Manipulating Cells and Viruses in Culture | 189 | Building and Fueling the Cell | 592 |
| 7 Recombinant DNA Technology | 221 | 14 Membrane Structure: The Plasma Membrane | 595 |
| 8 Genetic Analysis in Cell Biology | 263 | 15 Transport Across Cell Membranes | 633 |
| | | 16 Synthesis and Sorting of Plasma Membrane, Secretory, and Lysosomal Proteins | 669 |
| ► Part II | | 17 Cellular Genetics: Formation of ATP by Glycolysis and Oxidative Phosphorylation | 739 |
| Control of Cellular Activity by the Nucleus | 304 | 18 Photosynthesis | 779 |
| 9 The Molecular Anatomy of Genes and Chromosomes | 307 | | |

| | | | | | |
|--|---|-----|----|--|------|
| 19 | Organelle Biogenesis: The Mitochondrion, Chloroplast, Peroxisome, and Nucleus | 809 | 22 | Microfilaments: Cell Motility and Control of Cell Shape | 991 |
| ► Part IV | | | 23 | Microtubules and Intermediate Filaments | 1051 |
| Integrative and Specialized Cellular Events | | | 24 | Multicellularity: Cell-Cell and Cell-Matrix Interactions | 1123 |
| 20 | Cell-to-Cell Signaling | 850 | 25 | Regulation of the Eukaryotic Cell Cycle | 1201 |
| 21 | Nerve Cells | 853 | 26 | Cancer | 1247 |
| | | 925 | 27 | Immunity | 1295 |



| | | | |
|---|----------|---|----|
| <i>Chapter-Opening Illustrations</i> | xiv | RNA Is the Molecule That Is the Product of the Genome | 12 |
| ► Part I | | <i>Cells Have Both a Fixed Identity and an Ability to Change</i> | 12 |
| Laying the Groundwork | 1 | <i>Molecular Cell Biology</i> | 14 |
| 1 THE DYNAMIC CELL | 3 | 2 CHEMICAL FOUNDATIONS | 15 |
| <i>Evolution: Biology as a Historical Science</i> | 4 | <i>Energy</i> | 16 |
| <i>The Construction of Cells</i> | 5 | <i>Covalent Bonds</i> | 16 |
| Cells Are Surrounded by Water-Impermeable Membranes | 6 | Each Atom Can Make a Defined Number of Covalent Bonds | 17 |
| The Biological World Has Two Types of Cells | 7 | The Making or Breaking of Covalent Bonds Involves Large Energy Changes | 18 |
| Membranes Serve Functions Other Than Segregation | 8 | Covalent Bonds Exhibit Precise Orientations | 18 |
| Eukaryotic Cells Contain Organelles Thought to Have Evolved as Independent Organisms | 9 | Polar Covalent Bonds Result from Unequal Sharing of Electrons | 19 |
| <i>The Molecules of Life</i> | 9 | <i>Asymmetric Carbon Atoms and the Structure of Amino Acids and Carbohydrates</i> | 20 |
| Genetics Allowed DNA Organization to Be Analyzed Before the Structure of DNA Was Discovered | 10 | The α Carbon in Amino Acids Is a Chiral Carbon | 20 |
| The Ultimate Triumph of Genetics Is the Human Genome Project | 12 | Chiral Carbons Influence the Three-Dimensional Structure of Carbohydrates | 20 |

| | | | |
|--|----|--|----|
| <i>Noncovalent Bonds and the Structures of Biological Molecules</i> | 21 | Many Cellular Processes Involve the Transfer of Electrons in Oxidation-Reduction Reactions | 37 |
| The Hydrogen Bond Underlies Water's Biological Properties | 22 | An Unfavorable Chemical Reaction Can Proceed If It Is Coupled with an Energetically Favorable Reaction | 39 |
| Ionic Interactions Are Attractions Between Oppositely Charged Ions | 23 | Hydrolysis of the Phosphoanhydride Bonds in ATP Releases Substantial Free Energy | 40 |
| Van der Waals Interactions Are Caused by Transient Dipoles | 24 | ATP Is Used to Fuel Many Cellular Processes | 42 |
| Hydrophobic Bonds Cause Nonpolar Molecules to Adhere to Each Other | 25 | Polymers of Glucose with Specific Glycosidic Linkages Serve as Storage Reservoirs | 43 |
| Binding Specificity Can Be Conferred by Multiple Weak Bonds | 26 | <i>Activation Energy and Reaction Rate</i> | 44 |
| <i>Biomembranes: Hydrophobic Sheets Separating Aqueous Compartments</i> | 27 | Energy Is Required to Initiate a Reaction | 44 |
| Phospholipids Are the Principal Components of Biomembranes | 27 | Enzymes Catalyze Biochemical Reactions | 45 |
| Phospholipids Spontaneously Form Bilayers in Aqueous Solutions | 27 | <i>Summary</i> | 47 |
| <i>Chemical Equilibrium</i> | 28 | <i>Review Questions</i> | 48 |
| <i>pH and the Concentration of Hydrogen Ions</i> | 30 | <i>References</i> | 49 |
| Water Dissociates into Hydronium and Hydroxyl Ions | 30 | 3 PROTEIN STRUCTURE AND FUNCTION | 51 |
| Acids Release Hydrogen Ions, and Bases Combine with Hydrogen Ions | 31 | <i>General Structure of Proteins</i> | 52 |
| Biological Molecules Can Have Both Acidic and Basic Groups | 31 | Form and Function Are Inseparable in Protein Architecture | 52 |
| The Henderson-Hasselbalch Equation Describes the Relationships between pH and Equilibrium Constants for Acids and Bases | 32 | Amino Acids, the Building Blocks of Proteins, Differ Only in Their Side Chains | 52 |
| Buffers Maintain the pH of Cells and of Extracellular Fluids | 32 | Polypeptides Are Amino Acids Connected by Peptide Bonds | 56 |
| <i>Biochemical Energetics: Free Energy in Biochemical Reactions</i> | 34 | Four Levels of Structure Determine the Shape of Proteins | 56 |
| The Change in Free Energy, ΔG , Determines the Direction of a Chemical Reaction | 34 | Polypeptides Can Be Chemically Analyzed and Synthesized | 59 |
| The ΔG of a Reaction Depends on Changes in Heat and Entropy | 34 | Three-Dimensional Protein Structure Is Determined through X-Ray Crystallography and NMR Spectroscopy | 59 |
| Temperature, Concentrations of Reactants, and Other Parameters Affect the ΔG of a Reaction | 35 | Graphical Representations of Proteins Highlight Internal Organization or Surface Structures | 61 |
| The Standard Free-Energy Change, ΔG° , Can Be Determined from Measurement of the Equilibrium Constant, K_{eq} | 36 | Secondary Structures Are Crucial Elements in Protein Architecture | 63 |
| The Generation of a Concentration Gradient Requires an Expenditure of Energy | 37 | Motifs Are Regular Combinations of Secondary Structures | 66 |
| | | Structural and Functional Domains Are Modules of Tertiary Structure | 67 |
| | | Sequence Homology Suggests Functional and Evolutionary Relationships among Proteins | 68 |
| | | Many Proteins Contain Tightly Bound Prosthetic Groups | 70 |

| | | | |
|---|----|--|-----|
| Chemical Modifications Alter the Biological Activity of Proteins | 71 | <i>Summary</i> | 97 |
| Proteolytic Processing and Protein Splicing Alter Protein Activity | 72 | <i>Review Questions</i> | 99 |
| A Protein Can Be Unfolded by Heat, Extreme pH, and Certain Chemicals | 73 | <i>References</i> | 100 |
| Many Denatured Proteins Can Refold into Their Native State In Vitro | 73 | 4 NUCLEIC ACIDS, THE GENETIC CODE, AND PROTEIN SYNTHESIS | 101 |
| Folding of Proteins In Vivo Is Promoted by Chaperones | 74 | <i>Nucleic Acids: Linear Polymers of Nucleotides</i> | 102 |
| <i>Enzymes</i> | 75 | <i>DNA</i> | 103 |
| The Active Site of an Enzyme Is a Cage of Amino Acids That Binds Substrates and Catalyzes Reactions | 75 | The Native State of DNA Is a Double Helix of Two Antiparallel Chains with Complementary Nucleotide Sequences | 103 |
| Proteases Degrade Proteins by Reducing Activation Energy for Peptide-Bond Hydrolysis | 76 | The Two Strands Can Separate, Causing DNA to Denature | 108 |
| Coenzymes Are Essential for Certain Enzyme-Catalyzed Reactions | 78 | Many DNA Molecules Are Circular | 109 |
| Activity of Some Enzymes Depends on a Conformational Change Induced by Substrate Binding | 80 | Linking Number, Twist, and Writhe Describe DNA Superstructure | 110 |
| The Catalytic Activity of an Enzyme Can Be Characterized Mathematically | 81 | <i>RNA: The Basic Chemical Structure and Its Function in Gene Expression</i> | 111 |
| Enzymatic Activity Can Be Regulated by Various Mechanisms | 83 | <i>Rules for the Synthesis of Proteins and Nucleic Acids and Macromolecular Carpentry</i> | 113 |
| <i>Antibodies</i> | 86 | <i>Nucleic Acid Synthesis</i> | 114 |
| Antibodies Are Multidomain, Multisubunit Proteins | 86 | Nucleic Acid Polymerization Can Be Described by Four Rules | 115 |
| Antigen-Binding Site Complements the Surface of the Antigen | 87 | Organization of Genes in DNA Differs in Prokaryotes and Eukaryotes | 116 |
| Antibodies Are Valuable Tools for Identifying and Purifying Proteins | 87 | Eukaryotic Primary RNA Transcripts Are Processed to Form Functional mRNAs | 119 |
| Antibodies Can Catalyze Chemical Reactions | 87 | <i>Protein Synthesis: The Three Roles of RNA in Translation</i> | 119 |
| <i>Techniques for Purifying and Characterizing Proteins</i> | 88 | Messenger RNA Carries Information from DNA in a Three-Letter Genetic Code | 120 |
| Centrifugation Can Separate Particles and Molecules That Differ in Mass or Density | 88 | Experiments with Synthetic mRNAs and Trinucleotides Break the Genetic Code | 122 |
| Electrophoresis Separates Molecules According to Their Charge: Mass Ratio | 92 | Folded Structure of tRNA Is Integral to Its Function | 124 |
| Liquid Chromatography Resolves Proteins by Mass, Charge, and Binding Affinity | 94 | Aminoacyl-tRNA Synthetases Activate tRNA | 126 |
| Highly Specific Assays Detect Individual Proteins | 96 | Each tRNA Molecule Is Recognized by a Specific Aminoacyl-tRNA Synthetase | 128 |
| | | Ribosomes Are Protein-Synthesizing Machines | 128 |
| | | <i>The Steps in Protein Synthesis</i> | 133 |
| | | AUG Is the Initiation Signal in mRNA | 133 |

| | | | |
|--|-----|---|-----|
| Initiation Factors, tRNA, mRNA, and the Small Ribosomal Subunit Form an Initiation Complex | 133 | <i>Sorting Cells and Their Parts</i> | 161 |
| Ribosomes Provide Three tRNA-Binding (A, P, and E) during Protein Elongation | 136 | Flow Cytometry Is Used to Sort Cells Optically | 161 |
| Polypeptide Termination Requires Protein Factors That Specifically Recognize UAA, UAG, and UGA | 138 | Fractionation Methods Isolate Subcellular Structures | 161 |
| <i>Summary</i> | 138 | <i>The Biomembranes and Organelles of the Eukaryotic Cell</i> | 166 |
| <i>Review Questions</i> | 139 | The Plasma Membrane Has Many Varied and Essential Roles | 167 |
| <i>References</i> | 139 | The Eukaryotic Nucleus Is Bounded by a Double Membrane | 167 |
| 5 CELL ORGANIZATION, SUBCELLULAR STRUCTURE, AND CELL DIVISION | 141 | The Nucleus Contains the Nucleolus, a Fibrous Matrix, and DNA-Protein Complexes | 168 |
| <i>Prokaryotic and Eukaryotic Cells</i> | 142 | The Cytosol Contains Many Cytoskeletal Elements and Particles | 168 |
| Prokaryotes Have a Relatively Simple Structure | 142 | The Endoplasmic Reticulum Is an Interconnected Network of Internal Membranes | 170 |
| Eukaryotic Cells Have Complex Systems of Internal Membranes and Fibers | 143 | Golgi Vesicles Process Secretory and Membrane Proteins and Sort Them to Their Proper Destinations | 172 |
| Prokaryotes and Eukaryotes Contain Similar Macromolecules | 143 | Lysosomes Are Acidic Organelles That Contain a Battery of Degradative Enzymes | 173 |
| Prokaryotes and Eukaryotes Differ in the Amount of DNA per Cell | 144 | Vacuoles in Plant Cells Store Small Molecules and Enable the Cell to Elongate Rapidly | 174 |
| The Organization of DNA Differs in Prokaryotic and Eukaryotic Cells | 147 | Peroxisomes Produce and Degrade Hydrogen Peroxide | 174 |
| <i>Light Microscopy and Cell Architecture</i> | 148 | The Mitochondrion Is the Principal Site of ATP Production in Aerobic Cells | 174 |
| The Resolution of Standard Light (Bright-Field) Microscopy Is Limited to about 0.2 μm | 148 | Chloroplasts Are the Sites of Photosynthesis | 175 |
| Immunofluorescence Microscopy Reveals Specific Proteins and Organelles within a Cell | 150 | Cilia and Flagella Are Motile Extensions of the Eukaryotic Plasma Membrane | 175 |
| Fluorescence Microscopy Can also Measure the Local Concentration of Ca^{2+} Ions and the Intracellular pH | 153 | The Plasma Membrane Binds to the Cell Wall or the Extracellular Matrix | 176 |
| The Confocal Scanning Microscope Produces Vastly Improved Fluorescent Images | 154 | <i>Cell Division and the Cell Cycle</i> | 177 |
| Phase-Contrast and Nomarski Interference Microscopy Visualize Unstained Living Cells | 156 | In Prokaryotes DNA Replication Is Followed Immediately by Cell Division | 177 |
| <i>Electron Microscopy</i> | 158 | In Eukaryotic Cells DNA Synthesis and Cell Division Occur in Special Phases of the Cell Cycle | 177 |
| Transmission Electron Microscopy Depends on the Differential Scattering of a Beam of Electrons | 158 | Mitosis Is the Complex Process That Apportions the New Chromosomes Equally between Daughter Cells | 178 |
| Minute Details Can Be Visualized on Viruses and Subcellular Particles | 159 | Plant Cells Show Some Variations in Mitosis | 180 |
| Scanning Electron Microscopy Visualizes Details on the Surface of Cells or Particles | 161 | | |

| | | | |
|--|-----|---|-----|
| Initiation Factors, tRNA, mRNA, and the Small Ribosomal Subunit Form an Initiation Complex | 133 | <i>Sorting Cells and Their Parts</i> | 161 |
| Ribosomes Provide Three tRNA-Binding (A, P, and E) during Protein Elongation | 136 | Flow Cytometry Is Used to Sort Cells Optically | 161 |
| Polypeptide Termination Requires Protein Factors That Specifically Recognize UAA, UAG, and UGA | 138 | Fractionation Methods Isolate Subcellular Structures | 161 |
| <i>Summary</i> | 138 | <i>The Biomembranes and Organelles of the Eukaryotic Cell</i> | 166 |
| <i>Review Questions</i> | 139 | The Plasma Membrane Has Many Varied and Essential Roles | 167 |
| <i>References</i> | 139 | The Eukaryotic Nucleus Is Bounded by a Double Membrane | 167 |
| 5 CELL ORGANIZATION, SUBCELLULAR STRUCTURE, AND CELL DIVISION | 141 | The Nucleus Contains the Nucleolus, a Fibrous Matrix, and DNA-Protein Complexes | 168 |
| <i>Prokaryotic and Eukaryotic Cells</i> | 142 | The Cytosol Contains Many Cytoskeletal Elements and Particles | 168 |
| Prokaryotes Have a Relatively Simple Structure | 142 | The Endoplasmic Reticulum Is an Interconnected Network of Internal Membranes | 170 |
| Eukaryotic Cells Have Complex Systems of Internal Membranes and Fibers | 143 | Golgi Vesicles Process Secretory and Membrane Proteins and Sort Them to Their Proper Destinations | 172 |
| Prokaryotes and Eukaryotes Contain Similar Macromolecules | 143 | Lysosomes Are Acidic Organelles That Contain a Battery of Degradative Enzymes | 173 |
| Prokaryotes and Eukaryotes Differ in the Amount of DNA per Cell | 144 | Vacuoles in Plant Cells Store Small Molecules and Enable the Cell to Elongate Rapidly | 174 |
| The Organization of DNA Differs in Prokaryotic and Eukaryotic Cells | 147 | Peroxisomes Produce and Degrade Hydrogen Peroxide | 174 |
| <i>Light Microscopy and Cell Architecture</i> | 148 | The Mitochondrion Is the Principal Site of ATP Production in Aerobic Cells | 174 |
| The Resolution of Standard Light (Bright-Field) Microscopy Is Limited to about 0.2 μm | 148 | Chloroplasts Are the Sites of Photosynthesis | 175 |
| Immunofluorescence Microscopy Reveals Specific Proteins and Organelles within a Cell | 150 | Cilia and Flagella Are Motile Extensions of the Eukaryotic Plasma Membrane | 175 |
| Fluorescence Microscopy Can also Measure the Local Concentration of Ca^{2+} Ions and the Intracellular pH | 153 | The Plasma Membrane Binds to the Cell Wall or the Extracellular Matrix | 176 |
| The Confocal Scanning Microscope Produces Vastly Improved Fluorescent Images | 154 | <i>Cell Division and the Cell Cycle</i> | 177 |
| Phase-Contrast and Nomarski Interference Microscopy Visualize Unstained Living Cells | 156 | In Prokaryotes DNA Replication Is Followed Immediately by Cell Division | 177 |
| <i>Electron Microscopy</i> | 158 | In Eukaryotic Cells DNA Synthesis and Cell Division Occur in Special Phases of the Cell Cycle | 177 |
| Transmission Electron Microscopy Depends on the Differential Scattering of a Beam of Electrons | 158 | Mitosis Is the Complex Process That Apportions the New Chromosomes Equally between Daughter Cells | 178 |
| Minute Details Can Be Visualized on Viruses and Subcellular Particles | 159 | Plant Cells Show Some Variations in Mitosis | 180 |
| Scanning Electron Microscopy Visualizes Details on the Surface of Cells or Particles | 161 | | |

| | | | |
|--|-----|--|-----|
| Yeast Cells Have a Simplified Division | 181 | Six Animal-Virus Classes Are Recognized Based on Genome Composition and Pathway of mRNA Synthesis | 208 |
| Meiosis Is the Form of Cell Division in Which Haploid Germ Cells Are Produced from Diploid Cells | 181 | <i>Radioisotopes: Indispensable Tools for Following Biological Activity</i> | 212 |
| <i>Summary</i> | 185 | Several Factors Determine the Choice of a Radiolabel | 213 |
| <i>Review Questions</i> | 186 | Radiolabeled Molecules Can Be Detected by Visual and Quantitative Methods | 214 |
| <i>References</i> | 187 | Intracellular Precursor Pools Affect the Outcome of Pulse-Chase Experiments | 215 |
| 6 MANIPULATING CELLS AND VIRUSES IN CULTURE | 189 | Synthesis Time of Macromolecules Can Be Estimated from Labeling Experiments | 216 |
| <i>Growth of Microorganisms in Culture</i> | 190 | The Dintzis Experiment Demonstrated That Proteins Are Synthesized from the Amino End to the Carboxyl End | 216 |
| Many Microorganisms Can Be Grown in Minimal Medium | 190 | <i>Summary</i> | 218 |
| Mutant Strains of Bacteria and Yeast Can Be Isolated by Replica Plating | 190 | <i>Review Questions</i> | 219 |
| <i>Growth of Animal Cells in Culture</i> | 193 | <i>References</i> | 220 |
| Rich Media Are Required for Culture of Animal Cells | 193 | 7 RECOMBINANT DNA TECHNOLOGY | 221 |
| Most Cultured Animal Cells Only Grow on Special Solid Surfaces | 193 | <i>DNA Cloning with Plasmid Vectors</i> | 222 |
| Primary Cell Cultures Have a Finite Life Span | 194 | Plasmids Are Extrachromosomal Self-Replicating DNA Molecules | 222 |
| Transformed Cells Can Grow Indefinitely in Culture | 196 | <i>E. coli</i> Plasmids Can Be Engineered for Use as Cloning Vectors | 222 |
| <i>The Use of Hybrid Cells in Genetic Analysis of Animal Cells and Production of Monoclonal Antibody</i> | 198 | Plasmid Cloning Permits Isolation of DNA Fragments from Complex Mixtures | 224 |
| Genes Can Be Mapped to Specific Chromosomes with Interspecific Hybrid Cells | 199 | <i>Production of Recombinant Plasmids</i> | 225 |
| Mutants in Purine- and Pyrimidine-Salvage Pathways Are Good Selective Markers | 200 | Restriction Enzymes Cut DNA Molecules at Specific Sequences | 225 |
| Hybridomas Are Fused Lymphoid Cells That Make Monoclonal Antibodies | 201 | Many Restriction Enzymes Generate DNA Fragments with “Sticky” Ends | 226 |
| <i>Viruses: Structure and Function</i> | 202 | DNA Ligase Covalently Links Restriction Fragments | 227 |
| Viral Capsids Are Regular Arrays of One or a Few Types of Proteins | 202 | Restriction Fragments Are Readily Inserted into Plasmid Vectors | 228 |
| Most Viral Host Ranges Are Narrow | 204 | <i>Formation and Uses of Synthetic DNA</i> | 228 |
| Viruses Can Be Cloned and Counted in Plaque Assays | 205 | <i>λ-Phage Cloning Vectors and Construction of a Genomic Library</i> | 229 |
| Viral Growth Cycles Are Classified as Lytic or Lysogenic | 205 | Bacteriophage λ Can Be Modified for Use as a Cloning Vector and Assembled In Vitro | 230 |
| Four Types of Bacterial Viruses Are Widely Used in Biochemical and Genetic Research | 206 | Nearly Complete Genomic Libraries of Higher Organisms Can Be Prepared by λ Cloning | 231 |
| Experiments with Plant Viruses Proved That RNA Can Act as a Genetic Material | 208 | Larger DNA Fragments Can Be Cloned in Cosmids and Other Vectors | 233 |

| | | | |
|---|-----|---|-----|
| <i>Construction of a cDNA Library</i> | 234 | From Protein to Gene and from Gene to Protein with Recombinant DNA Technology | 256 |
| cDNAs Are Produced by Copying Isolated mRNAs with Reverse Transcriptase | 235 | <i>Summary</i> | 257 |
| cDNAs Can Be Enzymatically Converted to Double-Stranded DNA and Cloned | 235 | <i>Review Questions</i> | 258 |
| <i>Identification of Specific Clones in a Genomic or cDNA Library</i> | 236 | <i>References</i> | 260 |
| Membrane Hybridization Can Be Used to Screen a Library | 237 | 8 GENETIC ANALYSIS IN CELL BIOLOGY | 263 |
| Certain cDNAs and Synthetic Oligonucleotides Are Used as Probes | 238 | <i>The Isolation and Characterization of Mutants</i> | 264 |
| Expression Cloning Identifies Specific Clones Based on Properties of the Encoded Proteins | 240 | Mutations Are Recessive or Dominant | 264 |
| <i>Analyzing and Sequencing Cloned DNA</i> | 240 | Mutations Involve Large or Small DNA Alterations | 266 |
| Cleavage with an Appropriate Restriction Enzyme Separates a Cloned DNA from Its Vector | 240 | Mutations Occur Spontaneously and Can Be Induced | 267 |
| Gel Electrophoresis Resolves DNA Fragments of Different Size | 242 | Some Human Diseases Are Caused by Spontaneous Mutations | 268 |
| Multiple Restriction Sites Can Be Mapped on a Cloned DNA Fragment | 243 | Various Genetic Screens Are Used to Identify Mutants | 269 |
| Pulsed-Field Gel Electrophoresis Separates Large DNA Molecules | 244 | Complementation Analysis Determines If Different Mutations Are in the Same Gene | 274 |
| Nucleotide Sequencing of Cloned DNA Fragments Paves the Way for Sequencing Entire Genomes | 245 | Metabolic and Other Pathways Can Be Genetically Dissected | 274 |
| <i>Analysis of Specific Nucleic Acids in Complex Mixtures</i> | 248 | Suppressor Mutations Can Identify Genes Encoding Interacting Proteins | 274 |
| Southern Blotting Detects Specific DNA Fragments | 248 | <i>Genetic Mapping of Mutations</i> | 277 |
| Northern Blotting Detects Specific RNAs | 249 | Segregation Patterns Indicate Whether Mutations Are on the Same or Different Chromosomes | 277 |
| Nuclease Protection Is Used to Quantitate Specific RNAs and Map the DNA Regions Encoding Them | 249 | Chromosomal Mapping Locates Mutations on Particular Chromosomes | 277 |
| Transcription Start Sites Can Be Mapped by S1 Protection and Primer Extension | 251 | Recombinational Analysis Can Map Genes Relative to Each Other on a Chromosome | 279 |
| <i>Designing Expression Systems That Produce Abundant Amounts of Specific Proteins</i> | 252 | DNA Polymorphisms Are Used to Map Human Mutations | 279 |
| Full-Length Proteins Encoded by Cloned Genes Can Be Produced in <i>E. coli</i> Expression Systems | 252 | Some Chromosomal Abnormalities Can Be Mapped by Banding Analysis | 281 |
| Proteins with Post-Translational Modifications Can Be Produced in Eukaryotic Expression Systems | 253 | <i>Molecular Cloning of Genes Defined by Mutations</i> | 284 |
| Proteins Encoded by Cloned Genes and cDNAs Can Be Expressed In Vitro | 253 | Physical Maps of Human Chromosomes Y and 21 Have Been Constructed by Screening YAC Clones for Sequence-Tagged Sites | 284 |
| The Polymerase Chain Reaction: An Alternative to Cloning | 254 | Physical and Genetic Maps Can Be Correlated | 284 |
| | | Physical Mapping of Selected Genomic Regions is the First Step in Cloning Many Genes | 285 |

| | | | |
|--|-----|---|-----|
| Mutation-Defined Genes Are Identified in Candidate Regions by Comparing Mutant and Wild-Type DNA Structure and mRNA Expression | 287 | Pseudogenes Are Duplicated Genes That Have Become Nonfunctional | 314 |
| Protein Structure Is Deduced from cDNA Sequences | 289 | rRNAs, tRNAs, and Histones Are Encoded by Tandemly Repeated Genes | 315 |
| <i>Gene Replacement and Transgenic Animals</i> | 291 | <i>Discovery of Repetitious DNA Fractions</i> | 316 |
| Specific Sites in Cloned Genes Can Be Altered In Vitro | 291 | Repeated DNA Reassociates More Rapidly Than Nonrepeated DNA | 316 |
| DNA Can Be Transferred into Eukaryotic Cells in Various Ways | 292 | Reassociation Experiments Reveal Three Major Classes of Eukaryotic DNA | 317 |
| Normal Genes Can Be Replaced with Mutant Alleles in Yeast and Mice | 292 | <i>Simple-Sequence DNA</i> | 318 |
| Foreign Genes Can Be Introduced into Plants and Animals | 296 | Higher Eukaryotes Contain Several Types of Simple-Sequence DNA | 318 |
| Gene Therapy Involves Use of Transgenes to Treat Genetic Diseases | 299 | Most Simple-Sequence DNA Is Located in Specific Chromosomal Regions | 319 |
| <i>Summary</i> | 300 | Differences in Lengths of Simple-Sequence Tandem Arrays Permit DNA Fingerprinting | 319 |
| <i>Review Questions</i> | 301 | <i>Immediate-Repeat DNA and Mobile DNA Elements</i> | 320 |
| <i>References</i> | 302 | Movement of Bacterial Mobile Elements Is Mediated by DNA | 323 |

► **Part II**
Control of Cellular Activity by the Nucleus **304**

| | | | |
|--|-----|--|-----|
| 9 MOLECULAR ANATOMY OF GENES AND CHROMOSOMES | 307 | Two Major Categories of Retrotransposons Are Found in Eukaryotic Cells | 328 |
| <i>Molecular Definition of a Gene</i> | 308 | Most Mobile Elements in Yeast Are Viral Retrotransposons | 328 |
| Most Prokaryotic Genes Lack Introns and Those Encoding Related Proteins Form Operons, Which Produce Polycistronic mRNA | 308 | <i>Copia</i> Retrotransposons Are the Most Common <i>Drosophila</i> Mobile Elements | 333 |
| Most Eukaryotic Transcription Units Produce Monocistronic mRNAs | 308 | LINES and SINES, the Most Abundant Mobile Elements in Mammals, Are Nonviral Retrotransposons | 334 |
| Simple Eukaryotic Transcription Units Give Rise to One mRNA | 309 | Retrotransposed Copies of Cellular RNAs Are Present in Eukaryotic Chromosomes | 336 |
| Complex Eukaryotic Transcription Units Give Rise to Alternative mRNAs | 309 | <i>Functional Rearrangements in Chromosomal DNA</i> | 338 |
| Some Genes Do Not Encode Protein | 310 | Salmonella Flagellar Antigens Can Switch through Inversion of a Transcription-Control Region | 338 |
| <i>Organization of Genes on Chromosomes</i> | 311 | Yeast Mating Types Can Switch by Gene Conversion | 339 |
| Genomes of Higher Eukaryotes Contain Much "Nonfunctional" DNA | 311 | Trypanosome Surface Antigens Undergo Frequent Changes via Gene Conversion | 341 |
| Cellular DNA Content Does Not Correlate with Phylogeny | 312 | Generalized DNA Amplification Produces Polytene Chromosomes | 343 |
| About a Quarter to Half of All Eukaryotic Protein-Coding Genes Are Solitary | 313 | Localized DNA Amplification of rRNA and Other Genes Occurs in Some Eukaryotic Cells | 343 |
| Gene Families Are Formed by Gene Duplication and Encode Homologous Proteins | 313 | | |

| | | | |
|---|-----|--|-----|
| Vertebrate Genes Encoding Antibodies Are Assembled from Gene Segments by Controlled Deletion of Intervening DNA | 344 | DnaA Protein Initiates Replication in <i>E. coli</i> | 372 |
| <i>Organizing Cellular DNA into Chromosomes</i> | 344 | DnaB Is a Helicase That Unwinds Duplex DNA | 372 |
| Prokaryotic Chromosomes Contain Highly Compacted Circular DNA Molecules with a Single Replication Origin | 344 | Primase Catalyzes Formation of RNA Primers for DNA Synthesis | 374 |
| Eukaryotic Nuclear DNA Associates with Highly Conserved Histone Proteins to Form Chromatin | 346 | At a Growing Fork One Strand Is Synthesized Discontinuously from Multiple Primers | 374 |
| Chromatin Exists in Extended and Condensed Forms | 347 | DNA Polymerase III Synthesizes Both the Leading and Lagging Strands | 375 |
| <i>Morphology and Functional Characteristics of Eukaryotic Chromosomes</i> | 349 | The Leading and Lagging Strands Are Synthesized Concurrently | 376 |
| Chromosome Number and Shape Are Species Specific | 349 | Interaction of Tus Protein with Termination Sites Stops DNA Replication | 378 |
| Nonhistone Proteins Provide a Structural Scaffold for Long DNA Loops in Chromosomes | 349 | <i>Eukaryotic DNA Replication</i> | 378 |
| Chromatin Contains Small Amounts of DNA-Binding Proteins in Addition to Histones and Scaffold Proteins | 354 | Eukaryotic Proteins That Replicate SV40 DNA In Vitro Exhibit Similarities and Differences with <i>E. coli</i> Replication Proteins | 378 |
| Each Chromosome Contains One Linear DNA Molecule | 354 | Telomerase Prevents Progressive Shortening of Lagging Strands during DNA Replication | 380 |
| Stained Chromosomes Have Characteristic Banding Patterns | 354 | <i>Role of Topoisomerases in DNA Replication</i> | 381 |
| Heterochromatin Consists of Chromosome Regions That Do Not Uncoil | 355 | Type I Topoisomerases Relax DNA by Nicking and Closing One Strand of Duplex DNA | 381 |
| Three Functional Elements Are Required for Replication and Stable Inheritance of Chromosomes | 355 | Type II Topoisomerases Change DNA Topology by Breaking and Rejoining Double-Stranded DNA | 382 |
| Yeast Artificial Chromosomes Can Be Used to Clone Megabase DNA Fragments | 359 | Replicated Circular DNA Molecules Are Decatenated by Type II Topoisomerases | 383 |
| <i>Summary</i> | 359 | Linear Daughter Chromatids Also Are Separated by Type II Topoisomerases | 383 |
| <i>Review Questions</i> | 360 | <i>Repair of DNA</i> | 385 |
| <i>References</i> | 361 | Proofreading by DNA Polymerase Corrects Copying Errors | 385 |
| 10 DNA REPLICATION, REPAIR, AND RECOMBINATION | 365 | Environmental DNA Damage Can Be Repaired by Several Mechanisms | 386 |
| <i>General Features of Chromosomal Replication</i> | 366 | Excision Repair in <i>E. coli</i> Removes Bulky Chemical Adducts Caused by UV Light and Carcinogens | 387 |
| DNA Replication Is Semiconservative | 366 | Genetic Studies in Eukaryotes Have Identified DNA-Repair Genes | 388 |
| Most DNA Replication Is Bidirectional | 366 | <i>Recombination between Homologous DNA Sites</i> | 389 |
| DNA Replication Begins at Specific Chromosomal Sites | 370 | | |
| <i>DNA Replication in E. coli</i> | 372 | | |

| | | | |
|---|-----|---|-----|
| Holliday Recombination Model Is Supported by Observation of Predicted Intermediate Structures | 389 | Binding of <i>lac</i> Repressor to the <i>lac</i> Operator Blocks Transcription Initiation by RNA Polymerase | 416 |
| Recombination in <i>E. coli</i> Occurs by Three Similar Pathways That All Require RecA Protein | 391 | Most Bacterial Repressors Are Homodimers Containing α Helices That Insert into Adjacent Major Grooves of Operator DNA | 418 |
| Site-Specific Integration of λ Phage Mimics a Homologous Recombination Event | 395 | Positive Control of the <i>lac</i> Operon Is Exerted by cAMP-CAP | 421 |
| Studies in Yeast Are Providing Insights into Meiotic Recombination | 396 | Cooperative Binding of cAMP-CAP and RNA Polymerase to <i>lac</i> Control Region Activates Transcription | 423 |
| Gene Conversion Can Occur near the Crossover Point during Reciprocal Recombination | 396 | Control of Transcription from All Bacterial Promoters Involves Similar But Distinct Mechanisms | 423 |
| <i>Summary</i> | 400 | Transcription from Some Promoters Is Initiated by Alternative Sigma (σ) Factors | 424 |
| <i>Review Questions</i> | 401 | RNA Polymerase Containing σ^{54} Is Regulated by Proteins That Bind at Enhancer Sites Distant from the Transcription Initiation Site | 425 |
| <i>References</i> | 402 | <i>Eukaryotic Gene Control: Purposes and General Principles</i> | 426 |
| 11 REGULATION OF TRANSCRIPTION INITIATION | 405 | Most Genes in Higher Eukaryotes Are Regulated by Controlling Their Transcription | 427 |
| <i>Early Genetic Analysis of lac-Operon Control in E. coli</i> | 406 | DNA Regulatory Sites Often Are Located Many Kilobases from Eukaryotic Transcription Start Sites | 428 |
| Enzymes Encoded at <i>lac</i> Operon Can Be Induced and Repressed | 407 | <i>Structure and Function of Eukaryotic Nuclear RNA Polymerases</i> | 430 |
| Mutations in <i>lacI</i> Cause Constitutive Expression of the <i>lac</i> Operon | 407 | Three Eukaryotic Polymerases Catalyze Formation of Different RNAs | 430 |
| Operator Constitutive Mutations Identify Binding Site for <i>lac</i> Repressor | 408 | Eukaryotic RNA Polymerases Have Complex Subunit Structure | 430 |
| Mutations in Promoter Prevent Expression of <i>lac</i> Operon | 409 | The Largest Subunit in RNA Polymerase II Has an Essential Carboxyl-Terminal Repeat | 432 |
| Regulation of <i>lac</i> Operon Depends on Cis-Acting DNA Sequences and Trans-Acting Proteins | 409 | RNA Polymerase II Initiates Transcription at DNA Sequences Corresponding to the 5' Cap of mRNAs | 433 |
| <i>Molecular Mechanisms of Transcription Initiation in Bacteria</i> | 411 | <i>Cis-Acting Regulatory Sequences in Eukaryotic DNA</i> | 435 |
| Induction of the <i>lac</i> Operon Leads to Increased Synthesis of <i>lac</i> mRNA | 411 | The TATA Box Positions RNA Polymerase II for Transcription Initiation in Many Genes | 435 |
| <i>E. coli</i> RNA Polymerase Generally Initiates Transcription at a Unique Position on DNA Template | 411 | Promoter-Proximal Elements Help Regulate Many Eukaryotic Genes | 436 |
| Protein-Binding Sites in <i>lac</i> Control Region Have Been Identified by Sequence Comparison and Footprinting Experiments | 412 | | |
| RNA Polymerase Interacts with Specific Promoter Sequences | 413 | | |
| σ^{70} Subunit of RNA Polymerase Functions as an Initiation Factor | 414 | | |
| α -Subunit Dimer Bends to rRNA Promoters in -40 to -60 Region | 416 | | |

| | | | |
|--|-----|--|-----|
| Transcription by RNA Polymerase II Often Is Stimulated by Distant Enhancer Sites | 439 | Only Essential Function of Polymerase I Is to Produce pre-rRNA | 468 |
| Most Eukaryotic Genes Are Regulated by Multiple Transcription-Control Elements | 441 | Species-Specific Initiation Factors Are Utilized by RNA Polymerase I | 469 |
| <i>Eukaryotic Transcription Factors</i> | 442 | <i>Transcription by RNA Polymerase III</i> | 470 |
| Biochemical and Genetic Techniques Have Been Used to Identify Transcription Factors | 442 | tRNA Genes Bind Two Multisubunit Initiation Factors | 471 |
| Many Transcription Factors Are Modular Proteins Composed of Distinct Functional Domains | 445 | 5S-rRNA Gene Binds Three Initiation Factors | 471 |
| A Variety of Protein Structures Form the DNA-Binding Domains of Eukaryotic Transcription Factors | 447 | TBP Is Required for Transcription Initiation by All Three Eukaryotic RNA Polymerases | 472 |
| Heterodimeric Transcription Factors Increase Regulatory Diversity | 452 | <i>Other Transcription Systems</i> | 473 |
| A Diverse Group of Amino Acid Sequences Are Found in Activation Domains | 452 | T7 and Related Bacteriophages Express Monomeric, Largely Unregulated RNA Polymerases | 473 |
| <i>RNA Polymerase II Transcription-Initiation Complex</i> | 453 | Mitochondrial DNA Transcription Exhibits Features Typical of Bacteriophage, Bacteria, and the Eukaryotic Nucleus | 473 |
| Transcription-Initiation Complex Contains Many Proteins Assembled in a Specific Order | 453 | Chloroplasts Contain an RNA Polymerase Homologous to the <i>E. coli</i> Enzyme | 474 |
| Transcription Activators Influence Assembly of Initiation Complex | 456 | Archaeobacteria Have an RNA Polymerase and Putative General Transcription Factors Similar to Those of the Eukaryotic Nucleus | 475 |
| Some Eukaryotic Regulatory Proteins Function as Repressors | 456 | <i>Summary</i> | 475 |
| <i>Regulating the Activity of Eukaryotic Transcription Factors</i> | 456 | <i>Review Questions</i> | 478 |
| Expression of Many Transcription Factors Is Restricted to Specific Cell Types | 457 | <i>References</i> | 480 |
| Some Transcription Factors Are Controlled by Lipid-Soluble Hormones | 458 | 12 TRANSCRIPTION TERMINATION, RNA PROCESSING, AND POSTTRANSCRIPTIONAL CONTROL | 485 |
| Polypeptide Hormones Signal Phosphorylation of Some Transcription Factors | 462 | <i>Transcription Termination in Prokaryotes</i> | 486 |
| <i>Influence of Chromatin Structure on Eukaryotic Transcription Initiation</i> | 464 | Rho-Independent Termination Sites Have Characteristic Sequences | 486 |
| Association of Genes with Heterochromatin Can Lead to Their Repression | 464 | Attenuation Can Cause Premature Chain Termination | 486 |
| Transcriptionally Inactive DNA Regions Are Resistant to DNase I | 466 | Transcription Termination at Some Sites Requires Rho Factor | 488 |
| Cytosine Methylation Is Associated with Inactive Genes in Vertebrates | 468 | Antitermination Can Prevent Premature Chain Termination | 490 |
| <i>Transcription by RNA Polymerase I</i> | 468 | <i>Eukaryotic Transcription-Termination Control</i> | 491 |
| Pre-rRNA DNA from All Eukaryotes Is Similar | 468 | HIV Tat Protein Is an RNA-Binding Antitermination Protein | 492 |
| | | Premature Termination of <i>c-myc</i> Transcription Occurs in Nondividing Cells | 492 |

| | | | |
|---|-----|--|-----|
| RNA Polymerase II Pauses during Transcription of <i>Drosophila</i> Heat-Shock Genes under Normal Conditions | 492 | Stability of Cytoplasmic mRNAs Varies Widely | 522 |
| <i>mRNA Processing in Higher Eukaryotes</i> | 494 | Degradation Rate of Some Eukaryotic mRNAs Is Regulated | 524 |
| mRNA Precursors Are Associated with Abundant Nuclear Proteins Containing Conserved RNA-Binding Domains | 494 | Translation of a Few mRNAs Is Regulated by Specific RNA-Binding Proteins | 525 |
| HnRNP Proteins May Have Multiple Functions | 496 | Antisense RNA Regulates Translation of Transposase mRNA in Bacteria | 527 |
| Pre-mRNAs Are Cleaved at Specific 3' Sites and Rapidly Polyadenylated in Animal Cells | 498 | <i>Processing of rRNAs and tRNAs</i> | 528 |
| RNA-DNA Hybridization Reveals Spliced Out Introns | 498 | Pre-rRNA Binds Proteins, Then Is Cleaved and Methylated in the Nucleolus | 528 |
| Splice Sites in Pre-mRNAs Exhibit Short, Conserved Sequences | 500 | Pre-rRNA Genes Act as Nucleolar Organizers | 530 |
| Excision of Introns and Splicing of Exons in Pre-mRNA Occur via Two Transesterification Reactions | 501 | Self-Splicing Group I Introns in Some Pre-rRNAs Were the First Examples of Catalytic RNA | 530 |
| Small Nuclear Ribonucleoprotein Particles Assist in Splicing | 501 | Processing of Pre-tRNA Involves Cleavage, Modification of Bases, and Sometimes a Unique Type of Splicing | 532 |
| Portions of Two Different RNAs Are Trans-Spliced in Some Organisms | 508 | <i>RNA Editing</i> | 535 |
| Self-Splicing Group II Introns Provide Clues to Evolution of snRNPs | 508 | RNA Editing Regulates Protein Function in Mammals | 535 |
| Regulation of RNA Processing Controls Expression of Some Proteins | 509 | RNA Editing in Trypanosome Mitochondria Drastically Alters mRNA Sequences | 536 |
| <i>Subnuclear Organization and Transport of Nuclear mRNA to the Cytoplasm</i> | 513 | <i>Summary</i> | 537 |
| Most Transcription and RNA Processing Occurs in a Limited Number of Domains in Mammalian Cell Nuclei | 513 | <i>Review Questions</i> | 539 |
| Messenger Ribonucleoproteins (mRNPs) Exit the Nucleus through Nuclear Pore Complexes | 515 | <i>References</i> | 540 |
| 5'-Cap Structures Are Recognized by the Nuclear Transport Mechanism | 516 | 13 GENE CONTROL IN DEVELOPMENT | 543 |
| Pre-mRNAs Associated with Spliceosomes Are Not Transported to the Cytoplasm | 518 | <i>Lysogeny or Lysis in λ-Phage Infection of E. coli</i> | 544 |
| mRNA Remains Associated with Protein in the Nucleus and Cytoplasm | 518 | Phage Mutants Unable to Undergo Lysogeny Fall into Three Main Complementation Groups | 544 |
| Transport of mRNPs to the Cytoplasm Is Regulated by Some Viral Proteins | 518 | cI Protein Maintains Lysogeny by Repressing and Activating Transcription from Different Promoters | 545 |
| <i>Regulation of mRNA Cytoplasmic Localization, Stability, and Translation</i> | 520 | cII and cIII Proteins Are Critical to Establishment of Lysogeny | 547 |
| Some mRNAs Are Directed to Specific Cytoplasmic Sites by Sequences in Their 3' Untranslated Regions | 521 | Induction of Lytic Cycle Requires Derepression of <i>cro</i> Gene | 548 |
| | | Cro and cI Have Similar DNA-Binding Domains But Interact Differently with λ Operators | 549 |
| | | Choice between Lysis and Lysogeny Involves Regulatory Mechanisms Found in More Complex Developmental Systems | 550 |

| | | | |
|---|-----|---|------------|
| <i>Cell-Type Specification and Mating-Type Conversion in Yeast</i> | 550 | Specification of the Anterior Region Depends on the Maternal <i>bicoid</i> Gene | 568 |
| Cell Type-Specific Gene Expression in Yeast Is Regulated by Numerous DNA-Binding Proteins | 551 | Protein Encoded by Maternal <i>nanos</i> Gene Represses Translation of <i>hunchback</i> mRNA in Posterior Region | 572 |
| Mating-Type Conversion Is Determined by Transcriptional Regulation of the <i>HO</i> Locus | 554 | Hunchback Protein Regulates Expression of Several Gap Genes along the Anteroposterior Axis | 573 |
| Silencer Elements Repress Expression at <i>HML</i> and <i>HMR</i> | 556 | Initial Patterning along Dorsoventral Axis Depends on Dorsal Protein | 573 |
| <i>Myogenesis in Mammals</i> | 556 | Maternal Terminal Genes Regulate Early Patterning of the Extreme Anterior and Posterior Ends of the Embryo | 575 |
| Embryonic Somites Give Rise to Myoblasts, the Precursors of Skeletal Muscle Cells | 557 | Subsequent Anteroposterior Patterning Is Regulated by a Cascade of Transcription Factors Expressed from Three Groups of Zygotic Genes | 576 |
| Certain Fibroblasts Can Be Converted into Muscle (<i>Myotubes</i>) | 558 | Selector Genes Control Regional Identity and Development of Adult Structures | 581 |
| The <i>myoD</i> Gene Can Trigger Muscle Development | 558 | <i>Mammalian Homologs of Drosophila ANT-C and BX-C</i> | 584 |
| Myogenic Proteins Are HLH Transcription Factors | 560 | Mammalian Hox Genes Are Colinear Homologs of <i>Drosophila</i> HOM-C Genes | 584 |
| Myogenic Gene Activation Depends on Specific Amino Acids in MyoD | 560 | Mutations in Hox Genes Result in Homeotic Transformations in the Developing Mouse | 585 |
| Id Protein Inhibits Activity of MyoD | 560 | <i>Summary</i> | 587 |
| Knockout Experiments Have Demonstrated Role of Myogenic Proteins In Vivo | 560 | <i>Review Questions</i> | 588 |
| <i>Neurogenesis in Drosophila and Mice</i> | 561 | <i>References</i> | 590 |
| <i>Drosophila</i> Sensory Hairs Arise from Proneural Clusters, Which Express Achaete and Scute Proteins | 562 | | |
| A Single Sensory Organ Precursor Develops from a Proneural Cluster in <i>Drosophila</i> | 562 | | |
| <i>Drosophila</i> Neurogenesis and Mammalian Myogenesis May Occur via Analogous Pathways Involving HLH Proteins | 562 | | |
| MASH1, a Homolog of Achaete and Scute Proteins, Regulates Neurogenesis in the Mouse | 562 | | |
| Specification of Other Cell Types Is Controlled by Different Classes of Transcription Factors | 565 | | |
| <i>Regional Specification during Drosophila Embryogenesis</i> | 565 | | |
| <i>Drosophila</i> Has Two Life Forms | 565 | | |
| Patterning Information Is Generated during Oogenesis and Early Embryogenesis | 565 | | |
| Morphogens Regulate Development as a Function of Their Concentration | 568 | | |
| Four Maternal Gene Systems Regulate Regionalization in the Early Embryo | 568 | | |
| | | ► Part III | |
| | | Building and Fueling the Cell | 592 |
| | | 14 MEMBRANE STRUCTURE AND THE PLASMA MEMBRANE | 595 |
| | | <i>General Architecture of Lipid Membranes</i> | 596 |
| | | All Membranes Contain Phospholipids and Proteins | 596 |
| | | The Phospholipid Bilayer Is the Basic Structural Unit of Biological Membranes | 599 |
| | | Phospholipid Bilayers Exhibit Two-Dimensional Fluidity That Depends on Temperature and Composition | 599 |
| | | Several Types of Evidence Point to the Universality of the Phospholipid Bilayer | 602 |
| | | Phospholipid Bilayers and Biological Membranes Form Closed Compartments | 603 |

| | | | |
|--|-----|---|-----|
| <i>Membrane Proteins</i> | 604 | <i>Summary</i> | 628 |
| Proteins Interact with Membranes in Different Ways | 604 | <i>Review Questions</i> | 629 |
| Transmembrane Proteins Contain Long Segments of Hydrophobic Amino Acids Embedded in the Phospholipid Bilayer | 604 | <i>References</i> | 631 |
| Proteins Can Be Removed from Membranes by Detergents or High-Salt Solutions | 604 | 15 TRANSPORT ACROSS CELL MEMBRANES | 633 |
| Many Integral Proteins Contain Multiple Transmembrane α Helices | 606 | <i>Major Types of Membrane Transport Proteins</i> | 634 |
| Porins Are Transmembrane Proteins Composed of Multiple β Strands | 609 | <i>Diffusion of Small Molecules across Pure Phospholipid Bilayers</i> | 635 |
| Some Integral Proteins Are Bound to the Membrane by Covalently Attached Hydrocarbon Chains | 610 | <i>Uniporter-Catalyzed Transport of Specific Molecules</i> | 636 |
| Interfacial Catalysis Involves Soluble Enzymes Acting at Membrane Surfaces | 612 | Three Main Features Distinguish Uniport Transport from Passive Diffusion | 637 |
| The Orientation of Proteins in Membranes Can Be Experimentally Determined | 612 | Two General Models Have Been Proposed for Transporters | 638 |
| <i>Glycoproteins and Glycolipids</i> | 612 | Glucose Entry into Erythrocytes Is Mediated by a Uniporter | 638 |
| Many Integral Proteins Contain Sugars Covalently Linked to Their Exoplasmic Domains | 613 | <i>Ion Channels, Intracellular Ion Environment, and Membrane Electric Potential</i> | 640 |
| Many Glycolipids Are Located in the Cell-Surface Membrane | 614 | Ionic Gradients and Electric Potential Are Maintained across the Plasma Membrane | 640 |
| <i>Principles of Membrane Organization</i> | 615 | Certain K^+ Channels Generate the Membrane Electric Potential | 641 |
| All Integral Proteins Bind Asymmetrically to the Lipid Bilayer | 615 | Ion Concentration Gradients and Electric Potential Drive the Movement of Ions across Biological Membranes | 643 |
| The Two Membrane Leaflets Have Different Lipid Compositions | 616 | <i>Active Ion Transport and ATP Hydrolysis</i> | 644 |
| Freeze-Fracture and Deep-Etching Techniques Reveal the Two Membrane Faces in Electron Microscopy | 616 | Ion Pumps Can Be Grouped into Three Classes (P, V, and F) | 644 |
| Most Integral Proteins and Lipids Are Laterally Mobile in Biomembranes | 616 | Ca^{2+} ATPase Maintains Low Cytosolic Ca^{2+} Concentration | 645 |
| Some Membrane Proteins Interact with Cytoskeletal Components | 616 | Coupling of ATP Hydrolysis and Ion Pumping by P-Class ATPases Involves an Ordered Kinetic Mechanism | 647 |
| Erythrocytes Have an Unusual Plasma Membrane That Is Tightly Anchored to the Cytoskeleton | 620 | Na^+/K^+ ATPase Maintains the Intracellular Concentrations of Na^+ and K^+ in Animal Cells | 648 |
| <i>Specializations of the Plasma Membrane</i> | 623 | V-Class H^+ ATPases Pump Protons across Lysosomal and Vacuolar Membranes | 650 |
| Plasma Membranes of Polarized Cells Are Divided into Two Regions with Different Compositions and Functions | 623 | The Multidrug-Transport Protein is an ATP-Powered Pump and ATP-Dependent Cl^- Channel | 651 |
| Tight Junctions Seal Off Body Cavities and Restrict Diffusion of Membrane Components | 625 | <i>Cotransport Catalyzed by Symporters and Antiporters</i> | 652 |
| Desmosomes and Gap Junctions Interconnect Cells and Control Passage of Molecules between Them | 628 | Na^+ -Linked Symporters Import Amino Acids and Glucose into Many Animal Cells | 652 |

| | | | |
|--|-----|---|-----|
| Na ⁺ -Linked Antiporter Exports Ca ²⁺ from Cells | 654 | Phospholipids Move from the ER to Other Cellular Membranes | 673 |
| Band 3 Is an Anion Antiporter That Exchanges Cl ⁻ and HCO ₃ ⁻ across the Erythrocyte Membrane | 655 | <i>Sites of Synthesis of Organelle and Membrane Proteins</i> | 674 |
| H ⁺ /K ⁺ ATPase and Anion Antiporter Combine to Acidify the Stomach Contents While Maintaining Cytosolic pH Near Neutrality | 657 | All Nuclear-Encoded Proteins Are Made by the Same Cytosolic Ribosomes | 675 |
| Several Symporters and Antiporters Regulate Cytosolic pH | 658 | Membrane-Attached and Membrane-Unattached Ribosomes Synthesize Different Proteins | 676 |
| <i>Plant and Prokaryotic Membrane Transport Proteins</i> | 659 | <i>Overall Pathway for Synthesis of Secretory, Lysosomal, and Membrane Proteins</i> | 676 |
| H ⁺ Pumps and Anion Channels Establish Electric Potential and a Steep H ⁺ Concentration Gradient across the Plant-Vacuole Membrane | 659 | Newly Made Secretory Proteins Are Localized to the Lumen of the Rough ER | 676 |
| Proton Antiporters Enable Plant Vacuoles to Accumulate Metabolites and Ions | 659 | Many Organelles Participate in Protein Secretion | 677 |
| The Potential across the Plasma Membrane of Plant, Bacterial, and Fungal Cells Is Generated by Proton Pumping | 660 | All Secretory Proteins Move from the Rough ER to Golgi Vesicles to Secretory Vesicles | 677 |
| Proton Symporters Import Many Nutrients into Bacteria | 660 | The Steps in Protein Secretion Can Be Studied Genetically | 678 |
| <i>Osmosis, Water Channels, and the Regulation of Cell Volume</i> | 661 | Plasma Membrane Glycoproteins Follow the Same Maturation Pathway as Continuously Secreted Proteins | 680 |
| Osmotic Pressure Causes Water Movement across Membranes | 662 | <i>The Transport of Secretory and Membrane Proteins into or across the ER Membrane</i> | 681 |
| Water Channels Are Necessary for Bulk Osmotic Flow of Water across Membranes | 663 | A Signal Sequence on Nascent Secretory Proteins Targets Them to the ER and Is then Cleaved Off | 682 |
| Some Animal Cells Regulate Their Volume by Modulating Their Internal Osmotic Strength | 663 | Several Receptor Proteins Mediate the Interaction of Signal Sequences with the ER Membrane | 683 |
| Changes in Intracellular Osmotic Pressure Cause Leaf Stomata to Open | 664 | Polypeptides Cross the ER Membrane in Protein-Lined Channels | 686 |
| <i>Summary</i> | 665 | ATP-Hydrolyzing Chaperone Proteins Prevent Protein Misfolding and Are Essential for Translocation of Secretory Proteins into the ER | 687 |
| <i>Review Questions</i> | 666 | Topogenic Sequences in Integral Membrane Proteins Allow Them to Achieve Their Proper Orientation in the ER Membrane | 688 |
| <i>References</i> | 667 | <i>Post-Translational Modifications of Secretory and Membrane Proteins in the Rough ER</i> | 694 |
| 16 SYNTHESIS AND SORTING OF PLASMA MEMBRANE, SECRETORY, AND LYSOSOMAL PROTEINS | 669 | Disulfide Bonds Are Formed in the ER Lumen Soon after Synthesis | 694 |
| <i>The Synthesis of Membrane Lipids</i> | 671 | Chaperone Proteins Facilitate the Folding of Newly Made Proteins | 695 |
| Phospholipids Are Synthesized in Association with Membranes | 671 | | |
| Special Membrane Proteins Allow Phospholipids to Equilibrate in Both Membrane Leaflets | 671 | | |

| | | | |
|--|-----|---|-----|
| The Formation of Oligomeric Proteins Occurs in the ER | 696 | A Type of Coated Vesicle without Clathrin Mediates ER-to-Golgi Transport and Transport within the Golgi | 713 |
| <i>Quality Control in the ER</i> | 697 | The Steps in Vesicular Transport Can Be Studied Biochemically and Genetically | 714 |
| Only Properly Folded Proteins Are Transported from the Rough ER to the Golgi Complex | 697 | A Family of Small GTP-Binding Proteins May Target Transport Vesicles to Their Correct Destinations | 716 |
| Unassembled or Misfolded Proteins Are Often Degraded within the ER | 698 | <i>Golgi and Post-Golgi Sorting and Processing of Membrane and Secretory Proteins</i> | 718 |
| ER-Specific Proteins Are Retained in the Rough ER or Are Returned There from the <i>Cis</i> -Golgi | 698 | Sequences in the Membrane-Spanning Domain Cause the Retention of Proteins in the Golgi | 719 |
| <i>Protein Glycosylation: Discrete Steps in the ER and Golgi Complex</i> | 699 | Different Vesicles Are Used for Continuous and Regulated Protein Secretion | 719 |
| Different Structures Characterize <i>N</i> - and <i>O</i> -Linked Oligosaccharides | 699 | Secretory and Membrane Proteins Undergo Several Proteolytic Cleavages During the Late Maturation Stages | 720 |
| Nucleotide Sugars Are the Precursors of Oligosaccharides | 700 | The Proteolytic Maturation of Insulin Occurs in Acidic, Clathrin-Coated Secretory Vesicles | 722 |
| <i>O</i> -Linked Oligosaccharides Are Formed by the Sequential Addition of Sugars | 703 | <i>Sorting of Membrane Proteins Internalized from the Cell Surface</i> | 722 |
| The ER and Golgi Membranes Contain Transporters for Nucleotide Sugars | 703 | In Receptor-Mediated Endocytosis, Cell Surface Receptors Are Internalized in Clathrin-Coated Vesicles | 722 |
| The Diverse <i>N</i> -Linked Oligosaccharides Share Certain Structural Features That Reflect a Common Precursor | 704 | The Low-Density Lipoprotein (LDL) Receptor Binds and Internalizes Cholesterol-Containing Particles | 724 |
| The Processing <i>N</i> -Linked Oligosaccharides Involves the Sequential Removal and Addition of Sugar Residues | 704 | Mutant LDL Receptors Reveal a Signal for Internalizing Receptors into Clathrin-Coated Pits | 724 |
| Modifications to <i>N</i> -Linked Oligosaccharides Are Completed in the Golgi Vesicles | 706 | Receptors and Ligands Dissociate in an Acidic Late Endosome/CURL Organelle | 726 |
| The Movement of Proteins through the Secretory Pathway Can Be Monitored by Following the Processing of <i>N</i> -Linked Oligosaccharides | 707 | Transferrin Delivers Iron to Cells by Receptor-Mediated Endocytosis | 727 |
| <i>N</i> -Linked and <i>O</i> -Linked Oligosaccharides May Stabilize Maturing Secretory and Membrane Proteins | 708 | Some Proteins Internalized by Endocytosis Remain within the Cell, or Are Transported across the Cell and Secreted | 728 |
| Phosphorylated Mannose Residues Target Proteins to Lysosomes | 709 | Proteins Are Sorted in Several Different Ways to Different Domains of the Plasma Membrane | 729 |
| Genetic Defects Have Elucidated the Role of Mannose Phosphorylation | 711 | Viruses and Toxins Enter Cells by Receptor-Mediated Endocytosis | 731 |
| <i>The Mechanism and Regulation of Vesicular Transport to and from the ER and the Golgi Complex</i> | 711 | <i>Summary</i> | 734 |
| Two Types of Coated Vesicles Transport Proteins from Organelle to Organelle | 711 | <i>Review Questions</i> | 735 |
| Clathrin Forms a Lattice Shell around Coated Pits and Vesicles | 711 | <i>References</i> | 737 |
| A Chaperone Protein Catalyzes the Depolymerization of Clathrin-Coated Vesicles | 713 | | |

| | | | |
|--|-----|--|-----|
| 17 CELLULAR ENERGETICS: FORMATION OF ATP BY GLYCOLYSIS AND OXIDATIVE PHOSPHORYLATION | 739 | Most Electron Carriers Are Oriented in the Transport Chain in the Order of Their Reduction Potentials | 765 |
| <i>Energy Metabolism in the Cytosol</i> | 740 | Three Electron Transport Complexes Are Sites of Proton Translocation | 766 |
| In Glycolysis, ATP Is Generated by Substrate-Level Phosphorylation | 742 | The Q Cycle Increases the Number of Protons Transported by the CoQH ₂ -Cytochrome C Reductase Complex | 767 |
| Some Eukaryotic and Prokaryotic Cells Metabolize Glucose Anaerobically | 744 | The Cytochrome C Oxidase Complex Couples the Reduction of Oxygen to the Translocation of Protons | 768 |
| <i>Mitochondria and the Metabolism of Carbohydrates and Lipids</i> | 745 | <i>Metabolic Regulation</i> | 770 |
| The Outer and Inner Membranes of the Mitochondrion Are Structurally and Functionally Distinct | 745 | Respiration Is Controlled by the Production of ATP through the Proton-Motive Force | 770 |
| Acetyl CoA Is a Key Intermediate in the Mitochondrial Metabolism of Pyruvate and Fatty Acids | 748 | An Endogenous Uncoupler in Brown-Fat Mitochondria Converts H ⁺ Gradients to Heat | 770 |
| The Citric Acid Cycle Oxidizes the Acetyl Group of Acetyl CoA to CO ₂ and Reduces NAD and FAD to NADH and FADH ₂ | 749 | The Rate of Glycolysis Depends on the Cell's Need for ATP and Is Controlled by Multiple Allosteric Effectors | 771 |
| Electrons Are Transferred from NADH and FADH ₂ to Molecular O ₂ by Electron-Carrier Proteins | 750 | The Oxidation of Fatty Acids Occurs in Peroxisomes without Production of ATP | 772 |
| A Similar Electrochemical Protein Gradient Is Used to Generate ATP from ADP and P _i in Mitochondria, Bacteria, and Chloroplasts | 751 | <i>Summary</i> | 773 |
| <i>The Proton-Motive Force, ATP Generation, and Transport of Metabolites</i> | 752 | <i>Review Questions</i> | 774 |
| Closed Vesicles Are Required for the Generation of ATP | 752 | <i>References</i> | 776 |
| The Proton-Motive Force Is Composed of a Proton Concentration Gradient and a Membrane Electric Potential | 753 | 18 PHOTOSYNTHESIS | 779 |
| The F ₀ F ₁ Complex Couples ATP Synthesis to Proton Movement Down the Electrochemical Gradient | 753 | <i>An Overview of Photosynthesis</i> | 780 |
| Reconstitution of Close Membrane Vesicles Supports the Role of the Proton-Motive Force in ATP Synthesis | 756 | Photosynthesis Occurs on Thylakoid Membranes | 780 |
| Many Transporters in the Inner Mitochondrial Membrane Are Powered by the Proton-Motive Force | 758 | Photosynthesis Consists of Both "Light" and "Dark" Reactions | 782 |
| Inner-Membrane Proteins Allow the Uptake of Electrons from Cytosolic NADH | 759 | <i>The Light-Absorbing Step of Photosynthesis</i> | 783 |
| <i>NADH, Electron Transport, and Proton Translocation</i> | 759 | Each Photon of Light Has a Defined Amount of Energy | 783 |
| Electron Transport in Mitochondria Is Coupled to Proton Translocation | 759 | Chlorophyll <i>a</i> Is the Primary Light-Absorbing Pigment | 783 |
| The Mitochondrial Electron Transport Chain Transfers Electrons from NADH to O ₂ | 761 | The Absorption of Light by Reaction-Center Chlorophylls Causes a Charge Separation across the Thylakoid Membrane | 784 |
| | | <i>Molecular Analysis of Bacterial Photosynthesis</i> | 786 |
| | | Purple Photosynthetic Bacteria Utilize Only One Photosystem and Do Not Evolve O ₂ | 786 |

| | | | |
|--|-----|--|-----|
| Photoelectron Transport in the Photosynthetic Reaction Center of Purple Bacteria Results in a Charge Separation | 788 | Proteins Encoded by Mitochondria DNA Are Synthesized on Mitochondrial Ribosomes | 816 |
| Photosynthetic Bacteria also Carry Out Noncyclic Electron Transport | 790 | Mitochondrial Genetic Codes Are Different from the Standard Nuclear Code, and They Differ among Organisms | 816 |
| <i>Molecular Analysis of Photosynthesis in Plants</i> | 790 | In Animals, Mitochondrial RNAs Undergo Extensive Processing | 817 |
| Plants Utilize Two Photosystems, PSI and PSII, with Different Functions in Photosynthesis | 790 | Mutations in Mitochondrial DNA Cause Several Genetic Diseases in Man | 817 |
| Both PSI and PSII Are Essential for Photosynthesis in Chloroplasts | 792 | <i>Synthesis and Localization of Mitochondrial Proteins</i> | 819 |
| PSII Splits H ₂ O | 793 | Most Mitochondrial Proteins Are Synthesized in the Cytosol as Precursors | 819 |
| Electrons Are Transported from PSII to PSI | 794 | Matrix-Targeting Sequences Direct Imported Proteins to the Mitochondrial Matrix | 820 |
| PSI Forms NADPH | 796 | Mitochondrial Receptors Bind Matrix-Targeting Sequences | 824 |
| PSI Can Also Function in Cyclic Electron Flow | 796 | Intermediates in Translocation of Proteins into the Mitochondrion Can Be Accumulated and Studied | 824 |
| PSI and PSII Are Functionally Coupled | 796 | The Uptake of Mitochondrial Proteins Requires Energy | 824 |
| <i>CO₂ Metabolism during Photosynthesis</i> | 797 | Matrix Chaperones Are Essential for the Import and Folding of Mitochondrial Proteins | 826 |
| CO ₂ Fixation Is Catalyzed by Ribulose 1,5-Bisphosphate Carboxylase | 797 | Proteins Are Targeted to the Correct Submitochondrial Compartment by Multiple Signals and Several Pathways | 827 |
| CO ₂ Fixation Is Activated in the Light | 800 | Certain Mitochondrial Proteins Are Essential for Life | 829 |
| Photorespiration Liberates CO ₂ and Consumes O ₂ | 800 | The Synthesis of Mitochondrial Proteins Is Coordinated | 829 |
| Peroxisomes Play a Role in Photorespiration | 800 | <i>Chloroplast DNA and the Biogenesis of Chloroplasts and Other Plastids</i> | 830 |
| The C ₄ Pathway for CO ₂ Fixation Is Used by Several Tropical Plants | 802 | Chloroplast DNA Contains over 120 Different Genes | 830 |
| Sucrose Is Transported from Leaves through the Phloem to All Plant Tissues | 803 | Several Uptake-Targeting Sequences Direct Proteins Synthesized in the Cytosol to the Appropriate Chloroplast Compartment | 832 |
| <i>Summary</i> | 805 | Proplastids Can Differentiate into Chloroplasts or Other Plastids | 835 |
| <i>Review Questions</i> | 806 | <i>Peroxisome Biosynthesis</i> | 837 |
| <i>References</i> | 807 | All Peroxisomal Proteins Are Imported from the Cytosol | 837 |
| 19 ORGANELLE BIOGENESIS: THE MITOCHONDRION, CHLOROPLAST, PEROXISOME, AND NUCLEUS | 809 | Genetic Diseases Have Helped to Elucidate the Process of Peroxisome Biogenesis | 838 |
| <i>An Overview of Organelle Biogenesis Outside the Secretory Pathway</i> | 810 | | |
| <i>Mitochondrial DNA: Structure, Expression, and Variability</i> | 812 | | |
| Cytoplasmic Inheritance and DNA Sequencing Have Established the Existence of Mitochondrial Genes | 812 | | |
| The Size and Coding Capacity of mtDNA Varies in Different Organisms, Reflecting Evolutionary Movement of DNA between Mitochondrion and Nucleus | 813 | | |

| | | | |
|---|------------|--|-----|
| <i>Protein Traffic into and out of the Nucleus</i> | 840 | Analogous Regions in All Seven-spanning Receptors Determine G Protein and Ligand Specificity | 881 |
| Nuclear Proteins Are Selectively Imported into Nuclei | 840 | Degradation of cAMP Also Is Regulated | 881 |
| Nuclear Pores Are the Portals for Protein Transport | 841 | <i>Role of cAMP in the Regulation of Cellular Metabolism</i> | 881 |
| Multiple Types of Nuclear Localization Sequences Direct Proteins and Ribonucleoproteins to the Nucleus | 842 | cAMP and Other Second Messengers Activate Specific Protein Kinases | 881 |
| Receptor Proteins in Nuclear Pores Bind Nuclear Proteins for Import | 844 | Epinephrine Stimulates Glycogenolysis in Liver and Muscle Cells | 882 |
| <i>Summary</i> | 844 | cAMP-Dependent Protein Kinase Regulates the Enzymes of Glycogen Metabolism | 884 |
| <i>Review Questions</i> | 845 | Kinase Cascade Permits Multienzyme Regulation and Amplifies Hormone Signal | 885 |
| <i>References</i> | 847 | Cellular Responses to cAMP Vary among Different Cell Types | 885 |
| ► Part IV | | <i>Receptor Tyrosine Kinases</i> | 886 |
| Integrative and Specialized Cellular Events | 850 | SH2-Containing Proteins Bind to Specific Phosphotyrosine Residues in Activated RTKs | 886 |
| 20 CELL-TO-CELL SIGNALING: HORMONES AND RECEPTORS | 853 | Ras Protein Is a Key Component of RTK Signaling Pathways in Many Eukaryotes | 887 |
| <i>Overview of Extracellular Signaling</i> | 854 | Genetic Analysis of <i>Drosophila</i> Eye Development Identified Three Proteins That Link RTKs to a Kinase Cascade | 891 |
| Signaling Molecules Operate over Various Distances in Animals | 855 | GRB2 Is an Adapter Protein That Binds to Activated RTKs | 893 |
| Receptor Proteins Exhibit Ligand-Binding Specificity and Effector Specificity | 856 | Sos Protein Is Localized to the Plasma Membrane by Binding to the SH3 Domains in GRB2 | 893 |
| Hormones Can Be Classified Based on Their Solubility and Receptor Location | 856 | A Highly Conserved Kinase Cascade Transmits RTK-Mediated Signals Downstream from Ras | 894 |
| Effects of Many Hormones Are Mediated by Second Messengers | 857 | | |
| Cell-Surface Receptors Can Be Categorized into Four Major Classes | 859 | | |
| The Synthesis, Release, and Degradation of Hormones Are Regulated | 860 | | |
| <i>Identification and Purification of Cell-Surface Receptors</i> | 865 | | |
| Hormone Receptors Are Detected by Binding Assays | 865 | | |
| K_D Values for Cell-Surface Hormone Receptors Approximate the Concentrations of Circulating Hormones | 866 | | |
| Affinity Techniques Permit Purification of Receptor Proteins | 866 | | |
| Many Receptors Can Be Cloned without Prior Purification | 867 | | |
| <i>Seven-spanning G Protein-Linked Receptors</i> | 869 | | |
| Binding of Epinephrine to β - and α -Adrenergic Receptors Induces Tissue-Specific Responses Mediated by cAMP | 870 | | |
| | | Analogs Provide Information about Essential Features of Hormone Structure and Are Useful as Drugs | 871 |
| | | Studies with Mutant β -Adrenergic Receptors Identify Residues That Interact with Catecholamines | 872 |
| | | Trimeric Signal-Transducing G_s Protein Links β -Adrenergic Receptors and Adenylate Cyclase | 873 |
| | | $G_{s\alpha}$ Belongs to GTPase Superfamily of Intracellular Switch Proteins | 876 |
| | | Some Bacterial Toxins Irreversibly Modify G Proteins | 877 |
| | | Adenylate Cyclase Is Stimulated and Inhibited by Different Receptor-Ligand Complexes | 879 |

| | | | |
|---|-----|---|-----|
| Ras-Coupled RTKs Transduce Extracellular Signals by a Common Pathway | 896 | <i>Summary</i> | 918 |
| Yeast Mating-Factor Receptors Are Linked to G Proteins That Transmit Signals to MAP Kinase | 897 | <i>Review Questions</i> | 920 |
| <i>Other Important Second Messengers</i> | 899 | <i>References</i> | 922 |
| Cellular Effects of Ca ²⁺ Depend on Its Cytosolic Level and Often Are Mediated by Calmodulin | 899 | 21 NERVE CELLS | 925 |
| Ca ²⁺ Ions and cAMP Induce Hydrolysis of Muscle Glycogen | 901 | <i>Neurons, Synapses, and Nerve Circuits</i> | 926 |
| Inositol 1,4,5-Trisphosphate Causes the Release of Ca ²⁺ Ions from the ER | 901 | Specialized Regions of Neurons Carry Out Different Functions | 926 |
| Release of Intracellular Ca ²⁺ Stores Also Is Mediated by Ryanodine Receptors in Muscle Cells and Neurons | 904 | Synapses Are Specialized Sites Where Neurons Communicate with Other Cells | 929 |
| 1,2-Diacylglycerol Activates Protein Kinase C | 904 | Neurons Are Organized into Circuits | 931 |
| <i>Multiplex Signaling Pathways</i> | 905 | <i>The Action Potential and Conductance of Electric Impulses</i> | 932 |
| Some Activated RTKs Stimulate Activity of Phospholipase C _γ | 905 | The Resting Potential Is Generated Mainly by Open Potassium Channels | 933 |
| Multiple G Proteins Transduce Signals from Seven-Spanning Receptors to Different Effector Proteins | 905 | Opening and Closing Ion Channels Cause Specific, Predictable Changes in the Membrane Potential | 935 |
| G _{βγ} Acts Directly on Some Effectors in Mammalian Cells | 906 | Membrane Depolarizations Would Spread Only Short Distances without Voltage-Gated Cation Channels | 935 |
| <i>The Insulin Receptor and Regulation of Blood Glucose</i> | 907 | Opening of Voltage-Gated Sodium Channels Depolarizes the Nerve Membrane during Conductance of an Action Potential | 936 |
| Insulin Has Short-Term Effects on Glucose Metabolism and Long-Term Growth-Promoting Effects | 907 | Voltage-Dependent Sodium Channel Proteins Propagate Action Potentials Unidirectionally without Diminution | 938 |
| Insulin Signaling Pathway Involves a Soluble "Relay" Protein That Does Not Bind to the Receptor | 910 | Opening of Voltage-Gated Potassium Channels Causes Repolarization of the Plasma Membrane during an Action Potential | 938 |
| Insulin and Glucagon Work Together to Maintain a Stable Blood Glucose Level | 911 | Movements of Only a Few Sodium and Potassium Ions Generate the Action Potential | 939 |
| <i>Regulation of Cell-Surface Receptors</i> | 912 | Myelination Increases the Rate of Impulse Conduction | 940 |
| Receptors for Many Peptide Hormones Are Down-Regulated by Endocytosis | 912 | Action Potentials Are Generated in an All-or-Nothing Fashion by Summation of Electric Disturbances | 943 |
| Phosphorylation of Cell-Surface Receptors Modulates Their Activity | 913 | <i>Molecular Properties of Voltage-Gated Ion Channel Proteins</i> | 943 |
| <i>From Plasma Membrane to Nucleus</i> | 914 | Patch Clamps Permit Measurement of Ion Movements through Single Sodium and Potassium Channels | 944 |
| Activation of Some Transcription Factors Occurs via Several Signaling Pathways Coupled to G Protein-Linked Receptors and RTKs | 914 | All Voltage-Gated Ion Channels Have a Similar Molecular Structure | 946 |
| STATs Are Transcription Factors Activated by Protein Tyrosine Kinases Associated with Cell-Surface Receptors | 916 | | |

| | | | |
|--|-----|--|-----|
| <i>Shaker</i> Mutants in <i>Drosophila melanogaster</i> Led to the Cloning of a Large Family of Voltage-Gated Potassium Channel Proteins | 947 | Hydrolysis of Acetylcholine Terminates the Depolarization Signal | 965 |
| Study of Toxin-Resistant Mutants Led to the Identification of Amino Acids That Line the Ion-Conducting Pore of the Potassium Channel | 948 | <i>Functions of Other Neurotransmitters, Their Receptors, and Their Transporters</i> | 965 |
| A Complete <i>shaker</i> K ⁺ Channel Is Assembled from Four Subunits | 949 | GABA and Glycine Receptors Are Ligand-Gated Anion Channels Used at Many Inhibitory Synapses | 966 |
| The S4 Segment Comprises the Voltage Sensor | 949 | Cardiac Muscarinic Acetylcholine Receptor Activates a G Protein and Opens Potassium Channels | 967 |
| The N-terminal Segment of the <i>shaker</i> Protein Causes Channel Inactivation | 949 | Different Catecholamine Receptors Affect Different Intracellular Second Messengers | 967 |
| Potassium Channel Proteins Are Diverse | 950 | A Serotonin Receptor Modulates Potassium Channel Function via the Activation of Adenylate Cyclase | 968 |
| The Sodium Channel Protein Has Four Homologous Transmembrane Domains, Each Similar to a Potassium Channel Polypeptide | 951 | Neurotransmitter Transporters Are the Proteins Affected by Drugs Such as Cocaine | 970 |
| All Voltage-Gated Ion Channel Proteins Probably Evolved from a Common Ancestral Gene | 951 | Some Peptides Function as Both Neurotransmitters and Neurohormones | 970 |
| <i>Synapses and Impulse Transmission</i> | 952 | Endorphins and Enkephalins Are Neurohormones That Inhibit Transmission of Pain Impulses | 971 |
| Impulse Transmission across Electric Synapses Is Nearly Instantaneous | 952 | <i>Sensory Transduction: The Visual and Olfactory Systems</i> | 971 |
| Chemical Synapses Can Be Fast or Slow, Excitatory or Inhibitory, and Can Exhibit Signal Amplification and Computation | 953 | The Light-Triggered Closing of Sodium Channels Hyperpolarizes Rod Cells | 972 |
| Many Types of Receptors Bind the Same Neurotransmitter | 955 | Absorption of a Photon Triggers Isomerization of Retinal and Activation of Opsin | 974 |
| <i>Synaptic Transmission and the Nicotinic Acetylcholine Receptor</i> | 956 | Cyclic GMP Is a Key Transducing Molecule | 975 |
| Acetylcholine Is Synthesized in the Cytosol and Stored in Synaptic Vesicles | 956 | Rod Cells Adapt to Varying Levels of Ambient Light | 976 |
| Exocytosis of Synaptic Vesicles Is Triggered by Opening of Voltage-Gated Calcium Channels and a Rise in Cytosolic Calcium | 958 | Color Vision Utilizes Three Opsin Pigments | 977 |
| Multiple Proteins Align Synaptic Vesicles with the Plasma Membrane and Participate in Vesicle Exocytosis and Endocytosis | 960 | More Than a Thousand Different G-Protein-Coupled Receptors Detect Odors | 978 |
| The Nicotinic Acetylcholine Receptor Protein Is a Ligand-Gated Cation Channel | 962 | <i>Memory and Neurotransmitters</i> | 979 |
| Spontaneous Exocytosis of Synaptic Vesicles Produces Small Depolarizations in the Postsynaptic Membrane | 962 | Mutations in <i>Drosophila</i> Affect Learning and Memory | 979 |
| The Nicotinic Acetylcholine Receptor Contains Five Subunits, Each of Which Contributes to the Cation Channel | 963 | Gill-Withdrawal Reflex in <i>Aplysia</i> Exhibits Three Elementary Forms of Learning | 979 |
| | | A Novel Glutamate Receptor Is the Coincidence Detector in Long-Term Potentiation Exhibited by Many Synapses in the Mammalian Brain | 982 |

| | | | |
|---|------|--|------|
| Retrograde Signaling by the Gas Nitric Oxide May Be a Part of Long-Term Potentiation | 982 | A Family of Actin-Severing Proteins Generates New Filament Ends by Breaking Actin Filaments | 1007 |
| Mice Defective in the Hippocampal Ca ²⁺ -Calmodulin-Activated Protein Kinase Are Impaired in Long-Term Potentiation and in Spatial Learning—the Beginnings of a Molecular Psychology | 984 | Actin Filaments Are Stabilized by Actin-Capping Proteins | 1009 |
| <i>Summary</i> | 984 | Many Movements Are Driven by Actin Polymerization | 1010 |
| <i>Review Questions</i> | 986 | <i>Myosin: A Cellular Engine that Powers Motility</i> | 1012 |
| <i>References</i> | 987 | Myosin Is a Diverse Family of Proteins Characterized by Distinct Head, Neck, and Tail Domains | 1014 |
| 22 MICROFILAMENTS: CELL MOTILITY AND CONTROL OF CELL SHAPE | 991 | The Myosin Tail Domain Regulates Binding to Membranes or the Assembly of Thick Filaments | 1015 |
| <i>Actin Filaments</i> | 992 | The Myosin Head Domain Is an Actin-Activated ATPase | 1015 |
| All Eukaryotic Cells Contain Abundant Amounts of Actin | 994 | Myosin Heads Walk along Actin Filaments | 1015 |
| The Actin Sequence Has Changed Little during Evolution | 994 | A Myosin Head Takes an 11-nm Step Each Time an ATP Molecule Is Hydrolyzed | 1017 |
| ATP Holds Together the Two Lobes of the Actin Monomer | 994 | X-Ray Crystallography Reveals the Atomic Structure of the Motor Domain | 1018 |
| G-Actin Assembles into Long F-Actin Polymers | 995 | Conformational Changes in the Head Couple ATP Hydrolysis to Movement | 1020 |
| F-Actin Is a Helical Polymer of Identical Subunits | 995 | <i>Muscle, A Specialized Contractile Machine</i> | 1021 |
| F-Actin Has Structural and Functional Polarity | 996 | Some Muscles Contract, Others Generate Tension | 1022 |
| <i>Actin Architectures</i> | 996 | Striated Muscles Contain a Regular Array of Actin and Myosin | 1023 |
| The Actin Cytoskeleton Is Organized into Bundles and Networks of Filaments | 997 | In Smooth Muscle, Thick and Thin Filaments Are Not in Regular Arrays | 1025 |
| Actin Bundles and Networks Are Connected to the Membrane | 998 | Thick and Thin Filaments Slide Past Each Other During Contraction | 1025 |
| Cortical Networks of Actin Filaments Stiffen Cell Membranes and Immobilize Integral Membrane Proteins | 999 | A Third Filament System of Long Proteins Organizes the Sarcomere | 1026 |
| Dystrophin Anchors a Cortical Actin Network Directly to the Extracellular Matrix | 1000 | Calcium from the Sarcoplasmic Reticulum Triggers Contraction | 1026 |
| Actin Bundles Support Projecting Fingers of Membrane | 1002 | Calcium Activation of Myosin Light Chains Regulates Contraction in Smooth Muscle and Invertebrate Muscle | 1030 |
| <i>The Dynamics of Actin Assembly</i> | 1003 | <i>Actin and Myosin in Nonmuscle Cells</i> | 1032 |
| Actin Polymerization in Vitro Proceeds in Three Steps | 1003 | Actin and Myosin II Are Arranged in Sarcomere-Like Structures | 1032 |
| ATP Enhances Assembly from One End of a Filament | 1003 | Contractile Actin Bundles Are Attached to Specialized Sites at the Plasma Membrane | 1035 |
| Fungal Toxins Disrupt the Monomer-Polymer Equilibrium | 1006 | Myosin II Stiffens Cortical Membranes | 1036 |
| Actin-Binding Proteins Control the Lengths of Actin Filaments | 1006 | | |

| | | | |
|---|------|--|------|
| Actin and Myosin II Have Essential Roles in Cytokinesis | 1036 | <i>Kinesin, Dynein, and Intracellular Transport</i> | 1070 |
| Myosins I and V Move Membrane-Bounded Cargoes along Actin Filaments | 1038 | Fast Axonal Transport Occurs along Microtubules | 1070 |
| Membrane-Bound Myosin in Vesicle Movements | 1039 | Microtubules Provide Tracks for the Movement of Pigment Granules | 1072 |
| Myosin I and Myosin II Are Not Essentially Required for Cell Migration | 1039 | Intracellular Membrane Vesicles Travel Along Microtubules | 1072 |
| <i>Cell Motility</i> | 1040 | Microtubule Motor Proteins Promote Vesicle Translocation along Microtubules | 1075 |
| Movements of Fibroblasts Involve Controlled Polymerization and Rearrangements of Actin Filaments | 1041 | Kinesin Is a (+) End-Directed Motor Protein | 1075 |
| Ameboid Movement Involves Reversible Gel-Sol Transitions of an Actin Network | 1043 | Dynein Is a (-) End-Directed Motor Protein | 1078 |
| Cell Movements Are Coordinated by Various Second Messengers and Signal Transduction Pathways | 1044 | Multiple Motor Proteins Are Associated with Membrane Vesicles | 1078 |
| <i>Summary</i> | 1046 | <i>Cilia and Flagella: Structure and Movement</i> | 1079 |
| <i>Review Questions</i> | 1048 | All Eukaryotic Cilia and Flagella Contain Bundles of Doublet Microtubules | 1080 |
| <i>References</i> | 1049 | Ciliary and Flagellar Beating Is Produced by Controlled Sliding of Outer Doublet Microtubules | 1084 |
| 23 MICROTUBULES AND INTERMEDIATE FILAMENTS | 1051 | Dynein Arms Generate the Sliding Forces | 1084 |
| <i>Microtubule Structures</i> | 1052 | Axonemal Dyneins Are Multi-Headed Motor Proteins | 1086 |
| Tubulin Subunits Comprise the Wall of a Microtubule | 1052 | Flagellar Beat Requires Conversion of Sliding to Bending | 1086 |
| Microtubules Form a Diverse Array of Both Permanent and Transient Structures | 1054 | Genetic Studies Provide Information about the Roles of the Central Microtubules and the Radial Spokes | 1087 |
| Microtubules Grow from Microtubule-Organizing Centers | 1055 | Calcium Regulates the Direction of Swimming | 1087 |
| The Microtubule-Organizing Center Determines the Polarity of Cellular Microtubules | 1055 | Axonemes Assemble from Basal Bodies | 1088 |
| Multiple Tubulin Genes and Chemical Modification Leads to Tubulin Diversity | 1059 | Basal Bodies Closely Resemble Centrioles | 1089 |
| <i>Microtubule Dynamics</i> | 1061 | <i>Microtubule Dynamics and Motor Proteins during Mitosis</i> | 1090 |
| Microtubule Assembly and Disassembly Occur by Preferential Addition and Loss of $\alpha\beta$ Dimers at the (+) End | 1061 | The Mitotic Apparatus Is a Microtubule Machine for Separating Chromosomes | 1091 |
| Dynamic Instability Is an Intrinsic Property of Microtubules | 1064 | The Kinetochore Is a Specialized Attachment Site at the Chromosome Centromere | 1091 |
| Colchicine and Other Anti-Cancer Drugs Poison Microtubule Assembly or Disassembly | 1066 | Yeast Centromeres Bind a Single Microtubule | 1094 |
| <i>Microtubule-Associated Proteins</i> | 1067 | Centrosome Duplication and Migration During Interphase and Prophase Initiate the Assembly of the Mitotic Apparatus | |
| MAPs Organize Bundles of Microtubules | 1067 | | |
| MAPs Stabilize Microtubules | 1070 | | |

| | | | |
|---|------|---|------|
| During Prophase, Kinesin-Related Proteins and Cytoplasmic Dynein Participate in the Movements of Kinetochores and Centrosomes | 1097 | Collagen Is Assembled into Fibrils after Secretion | 1133 |
| Assembly of the Mitotic Apparatus Involves Dynamic Microtubules | 1098 | Mutations in Collagen Reveal Aspects of Its Structure and Biosynthesis | 1134 |
| At Metaphase Forces at the Kinetochores Move Chromosomes to the Equator of the Spindle | 1100 | Collagens Form Diverse Structures | 1134 |
| During Anaphase Chromosomes Separate and the Spindle Elongates | 1100 | Type IV Collagen Forms the Two-Dimensional Reticulum of the Basal Lamina | 1135 |
| Astral Microtubules Determine Where Cytokinesis Takes Place | 1103 | <i>Hyaluronan and Proteoglycans</i> | 1136 |
| Plant Cells Build a New Cell Wall During Cell Division | 1105 | Hyaluronan Is an Immensely Long, Negatively Charged Polysaccharide That Forms Hydrated Gels | 1136 |
| <i>Intermediate Filaments</i> | 1106 | Hyaluronan Inhibits Cell-Cell Adhesion and Facilitates Cell Migration | 1137 |
| Intermediate Filaments Are Classified into Five Types | 1106 | Proteoglycans Comprise a Diverse Family of Cell-surface and Extracellular-matrix Macromolecules | 1139 |
| All Subunit Proteins of Intermediate Filaments Have a Similar Structure | 1109 | Proteoglycans Can Bind Many Growth Factors | 1142 |
| Intermediate Filaments Are Dynamic Polymers in the Cell | 1111 | <i>Multiadhesive Matrix Proteins and Their Cell-Surface Receptors</i> | 1143 |
| Phosphorylation of the N-Terminal Domain Regulates Polymerization of Intermediate Filaments during Mitosis | 1112 | Laminin and Nidogen Are Principal Structural Proteins of All Basal Laminae | 1143 |
| Intermediate Filament-Associated Proteins Cross-Link Intermediate Filaments to Membranes and Microtubules | 1113 | Integrins Are Cell-Surface Receptors That Mediate Adhesion to the Extracellular Matrix and Cell-Cell Interactions | 1144 |
| <i>Summary</i> | 1116 | Fibronectins Bind Many Cells to Fibrous Collagens and Other Matrix Components | 1146 |
| <i>Review Questions</i> | 1118 | Fibronectins Promote Cell Adhesion to the Substratum | 1148 |
| <i>References</i> | 1119 | Fibronectins Promote Cell Migration | 1149 |
| 24 MULTICELLULARITY: CELL-CELL AND CELL-MATRIX INTERACTIONS | 1123 | <i>Cell-Cell Adhesion: Adhesive Proteins</i> | 1150 |
| <i>The Extracellular Matrix: Primary Components and Functions</i> | 1124 | Adhesive Proteins Mediate Cell-Cell Interactions | 1150 |
| <i>Collagen: A Class of Multifunctional Fibrous Proteins</i> | 1126 | E-Cadherin Is a Key Adhesive Protein Expressed by Epithelial Cells | 1150 |
| The Basic Structural Unit of Collagen Is a Triple Helix | 1127 | Cadherins Influence Morphogenesis and Differentiation | 1152 |
| Most Exons in Fibrous Collagen Genes Encode Gly-X-Y Sequences | 1128 | N-CAMs Mediate Ca ²⁺ -Independent Adhesion of Cells in Nervous Tissue and Muscle | 1153 |
| Collagen Fibrils Form by Lateral Interactions of Triple Helices | 1128 | Movement of Leukocytes into Tissues Requires Sequential Interaction of Specific Adhesive Proteins | 1153 |
| Denatured Collagen Polypeptides Cannot Renature to Form a Triple Helix | 1131 | <i>Cell-Cell Adhesion: Cell Junctions</i> | 1155 |
| Procollagen Chains Assemble into Triple Helices in the Rough ER and Are Modified in the Golgi Complex | 1131 | Three Types of Desmosomes Impart Rigidity to Tissues | 1156 |
| | | Intermediate Filaments Stabilize Epithelia by Connecting Spot Desmosomes | 1156 |

| | | | |
|--|------|--|------|
| Hemidesmosomes Connect Epithelial Cells to the Basal Lamina | 1157 | Extracellular Signals Can Repel Growth Cones | 1181 |
| Gap Junctions Allow Small Molecules to Pass between Adjacent Cells | 1157 | Different Growth Cones Navigate along Different Axons | 1182 |
| Connexin, a Transmembrane Protein, Forms Cylindrical Channels in Gap Junctions | 1159 | The Basal Lamina at the Neuromuscular Junction Directs Differentiation of Regenerating Nerve and Muscle | 1185 |
| <i>Dorsoventral Patterning During Embryogenesis</i> | 1159 | <i>Structure and Function of the Plant Cell Wall</i> | 1187 |
| Embryologic Development Is Directed by Induction | 1160 | Cellulose Molecules Form Long, Rigid Microfibrils | 1188 |
| Transforming Growth Factor β (TGF β) Has Numerous Inductive Effects in Invertebrates and Vertebrates | 1161 | Other Polysaccharides Bind to Cellulose to Generate a Complex Wall Matrix | 1188 |
| TGF β Homolog Encoded by the <i>decapentaplegic</i> Gene Controls Dorsoventral Patterning in <i>Drosophila</i> Embryos | 1162 | Cell Walls Contain Lignin and an Extended Hydroxyproline-Rich Glycoprotein | 1190 |
| Sequential Inductive Events Regulate Early <i>Xenopus</i> Development | 1163 | Plants Grow Primarily by Auxin-Induced Cell Enlargement | 1190 |
| <i>Formation of Internal Organs and Organization of Tissues</i> | 1167 | The Orientation of Newly Made Cellulose Microfibrils Is Affected by the Microtubule Network | 1192 |
| The Basal Lamina Is Essential for Differentiation of Many Epithelial Cells | 1167 | Plasmodesmata Interconnect the Cytoplasm of Adjacent Cells in Higher Plants | 1193 |
| Direct Cell-Cell Contact Regulates Kidney Induction | 1168 | <i>Summary</i> | 1194 |
| Hedgehog Organizes Pattern in the Chick Limb and the <i>Drosophila</i> Wing | 1169 | <i>Review Questions</i> | 1196 |
| <i>Developmental Regulation by Direct Cell-Cell Contact</i> | 1172 | <i>References</i> | 1197 |
| Boss Is a Cell-Surface Inductive Ligand for the Sev Receptor | 1172 | 25 REGULATION OF THE EUKARYOTIC CELL CYCLE | 1201 |
| Cell-Surface Notch and Delta Proteins Control Signaling between Many Different Types of Cells | 1173 | <i>Phases of the Cell Cycle</i> | 1202 |
| <i>Regulation of Neuronal Outgrowth</i> | 1175 | <i>Experimental Systems in Cell-Cycle Research</i> | 1203 |
| Individual Neurons Can Be Identified Reproducibly and Studied | 1175 | <i>Control of Entry into and Exit from Mitosis</i> | 1203 |
| Growth Cones Guide the Migration and Elongation of Developing Axons | 1176 | The Same Factor Promotes Oocyte Maturation and Mitosis in Somatic Cells | 1203 |
| Different Neurons Navigate along Different Outgrowth Pathways | 1177 | Mitotic Cycling in Early Embryos Depends on Synthesis and Degradation of Cyclin B | 1207 |
| Different Extracellular-Matrix Components Are Permissive for Neuronal Outgrowth | 1178 | MPF-Catalyzed Phosphorylation of Nuclear Lamins and Other Proteins Induces Early Mitotic Events | 1212 |
| Three Genes Define Dorsoventral Outgrowth in <i>C. elegans</i> | 1179 | Protein Degradation and Dephosphorylation Trigger Late Mitotic Stages | 1216 |
| A Chemoattractant Related to Unc6 Is Produced in the Floor Plate of the Vertebrate Neural Tube | 1180 | Biochemical Studies with <i>Xenopus</i> Egg Extracts Identified Central Role of MPF in Regulating Entry into Mitosis | 1219 |
| | | <i>Regulation of MPF Activity</i> | 1219 |
| | | MPF Catalytic Subunit Is Encoded by <i>cdc2+</i> Gene in <i>S. pombe</i> | 1221 |

| | | | |
|---|------|---|------|
| MPF Catalytic Subunit Contains Activating and Inhibitory Sites That Are Phosphorylated | 1222 | Alterations in Cell-to-Cell Interactions Are Associated with Malignancy | 1249 |
| Structure of Human Cyclin-Dependent Kinase 2 Suggests How Phosphorylation Regulates MPF Activity | 1224 | Tumor Cells Lack Normal Controls on Cell Growth | 1251 |
| Entry into Mitosis Is Controlled by Multiple Mechanisms That Regulate MPF Activity | 1224 | <i>Use of Cell Cultures in Cancer Research</i> | 1251 |
| <i>Control of Entry into the S Phase</i> | 1226 | Fibroblastic, Epithelial, and Nonadherent Cells Grow Readily in Culture | 1251 |
| <i>S. cerevisiae</i> Cdc28 Is Functionally Equivalent to <i>S. pombe</i> Cdc2 | 1227 | Some Cell Cultures Give Rise to Immortal Cell Lines | 1252 |
| S Phase–Promoting Factor Consists of a Catalytic Subunit and G ₁ Cyclin | 1228 | Certain Factors in Serum Are Required for Long-Term Growth of Cultured Cells | 1253 |
| Various Cdc28-Cyclin Heterodimers Regulate Progress through the Cell Cycle in <i>S. cerevisiae</i> | 1230 | Malignant Transformation Leads to Many Changes in Cultured Cells | 1254 |
| Cdc28-G ₁ Cyclin Complexes May Activate Transcription Factors at START | 1230 | Transcription of Oncogenes Can Trigger Transformation | 1258 |
| A Cdk-Cyclin Complex Regulates a DNA Initiation Factor in <i>Xenopus</i> | 1231 | <i>Oncogenes and Their Proteins: Classification and Characteristics</i> | 1258 |
| <i>Cell-Cycle Control in Mammalian Cells</i> | 1232 | Oncogenes Were Initially Identified in Viruses and Tumor Cell DNA | 1258 |
| Mammalian Restriction Point Is Analogous to START in Yeast Cells | 1232 | Five Types of Proteins Participate in Control of Cell Growth | 1259 |
| Multiple Cdks and Cyclins Regulate Passage of Mammalian Cells through the Cell Cycle | 1232 | Oncoproteins Affect the Cell's Growth-Control Systems in Various Ways | 1260 |
| Growth Factor–Induced Expression of Two Classes of Genes Returns G ₀ Mammalian Cells to the Cell Cycle | 1235 | Apoptosis, or Induced Cell Suicide, Is One Mechanism of Protection Against Cancer | 1267 |
| Activity of Transcription Factor E2F Is Required for Entry into S Phase | 1236 | Oncoproteins Act Cooperatively in Transformation and Tumor Induction | 1267 |
| <i>Role of Checkpoints in Cell-Cycle Regulation</i> | 1237 | Consistent Chromosomal Anomalies Associated with Tumors Point to the Presence of Oncogenes | 1268 |
| Presence of Unreplicated DNA Prevents Entry into Mitosis | 1238 | Inherited Human Propensities to Develop Cancer Point to Tumor-Suppressor Genes | 1269 |
| Defects in Assembly of the Mitotic Spindle Prevent Exit from Mitosis | 1239 | <i>DNA Viruses as Transforming Agents</i> | 1270 |
| DNA Damage Prevents Entry into S Phase and Mitosis | 1239 | DNA Viruses Can Transform Nonpermissive Cells by Random Integration of the Viral Genome into the Host-Cell Genome | 1270 |
| <i>Summary</i> | 1240 | Transformation by DNA Viruses Requires Interaction of a Few Independently Acting Viral Proteins | 1271 |
| <i>Review Questions</i> | 1242 | <i>RNA-Containing Retroviruses as Transforming Agents</i> | 1273 |
| <i>References</i> | 1243 | Virion-Producing Infection Cycle of Retroviruses Requires Integration into the Host-Cell Genome | 1273 |
| 26 CANCER | 1247 | Oncogenic Transducing Retroviruses Convert Cellular Proto-Oncogenes into Oncogenes | 1275 |
| <i>Characteristics of Tumor Cells</i> | 1248 | | |
| Malignant Tumor Cells Are Invasive and Can Spread | 1248 | | |

| | | | |
|--|------|---|------|
| Nononcogenic Transducing Retroviruses Have Been Constructed Experimentally | 1277 | The Immune System Has a Memory | 1304 |
| Slow-Acting Carcinogenic Retroviruses Can Activate Nearby Cellular Proto-Oncogenes after Integration into the Host-Cell Genome | 1278 | Other Parts of the Immune Response Are Carried Out by T Lymphocytes | 1306 |
| <i>Human Tumor Viruses</i> | 1279 | B Cells and T Cells Have Identifying Surface Markers | 1306 |
| <i>Chemical Carcinogens</i> | 1280 | Macrophages Play a Central Role in Stimulating Immune Responses | 1307 |
| Most Chemical Carcinogens Must Undergo Metabolic Conversion to Become Active | 1280 | Cells Responsible for the Immune Response Circulate throughout the Body | 1308 |
| The Carcinogenic Effect of Chemicals Depends on Their Interaction with DNA | 1284 | Tolerance Is a Central Concept of Immunology | 1309 |
| <i>The Role of Radiation and DNA Repair in Carcinogenesis</i> | 1284 | Immunopathology Is Disease Caused by the Immune System | 1309 |
| Ineffective or Error-Prone Repair of Damaged DNA Perpetuates Mutations | 1285 | <i>Antibodies, B Cells, and the Generation of Diversity</i> | 1310 |
| Some Defects in DNA-Repair Systems Are Associated with High Cancer Rates in Humans | 1285 | Heavy-Chain Structure Differentiates the Classes of Antibodies | 1310 |
| <i>The Multicausal, Multistep Nature of Carcinogenesis</i> | 1286 | Antibodies Have a Domain Structure | 1310 |
| Epigenetic Alterations May Occur in Teratocarcinomas | 1286 | The N-Terminal Domains of H and L Chains Have Highly Variable Structures That Constitute the Antigen-Binding Site | 1311 |
| Some Cancer-Inducing Chemicals Act Synergistically | 1286 | Several Mechanisms Generate Antibody Diversity | 1315 |
| Natural Cancers Result from the Interaction of Multiple Events over Time | 1287 | DNA Rearrangement Generates Antibody Diversity | 1315 |
| <i>Human Cancer</i> | 1287 | A Single Recombination Event Generates Diversity in L Chains | 1315 |
| <i>Summary</i> | 1288 | Imprecision of Joining Makes an Important Contribution to Diversity | 1318 |
| <i>Review Questions</i> | 1289 | Lambda Diversity Derives from Multiple Constant Regions | 1319 |
| <i>References</i> | 1291 | H-Chain Variable Regions Derive from Three Libraries | 1319 |
| 27 IMMUNITY | 1295 | Recognition Sequences for All Joining Reactions Are Virtually Indistinguishable | 1320 |
| <i>Overview</i> | 1296 | The Synthesis of Immunoglobulins Is Like That of Other Extracellular Proteins | 1321 |
| Antibodies Bind to Epitopes and Have Two Functional Domains | 1297 | <i>The Antigen-Independent Phase of B-Lymphocyte Maturation</i> | 1321 |
| Antibody Reaction with Antigen Is Reversible | 1299 | B-Lymphocytes Go through an Orderly Process of Gene Rearrangement | 1321 |
| Antibodies Come in Many Classes | 1300 | The Antigen-Independent Phase Can Generate Cells with 10^{11} Different Specificities | 1322 |
| Antibodies Are Made by B Lymphocytes | 1302 | | |
| The Immune System Has Extraordinary Versatility | 1302 | | |
| Clonal Selection Theory Underlies All Modern Immunology | 1303 | | |

| | | | |
|---|------|--|------|
| The Immune System Requires Allelic Exclusion | 1323 | Antibody Secretion by Activated B Cells Entails Many Cellular Changes | 1335 |
| Antibody Gene Rearrangement and Expression Are Controlled by Transcription Factors | 1323 | The Activation Process Produces Both Plasma Cells and Memory Cells | 1335 |
| <i>T Lymphocytes</i> | 1324 | The Change from Surface Antibody to Secreted Antibody Requires the Synthesis of an Altered H Chain | 1335 |
| There Are Two T-Cell Receptor Molecules | 1325 | Antibody Class Switching Also Requires an Altered H Chain | 1337 |
| T-Cell Receptors Recognize a Foreign Antigen in Combination with a Self-Molecule | 1325 | Memory Cells Participate in the Secondary Immune Response | 1338 |
| T Cells Are Educated in the Thymus to React with Foreign Proteins but Not Self-Proteins | 1329 | Somatic Mutation of Variable Regions Follows from Activation | 1338 |
| T Cells Respond to Antigen by Either Killing Cells or Secreting Protein Factors | 1330 | Tolerance Is Achieved Partly by Making B Cells Unresponsive | 1339 |
| <i>The Antigen-Dependent Phase of the Immune Response</i> | 1332 | <i>Summary</i> | 1340 |
| B-Cell Activation Involves B Cell–T _H Cell Collaboration | 1333 | <i>Review Questions</i> | 1341 |
| Activation Is the Result of Cellular Stimulation | 1335 | <i>References</i> | 1342 |
| | | <i>Glossary</i> | G-1 |
| | | <i>Index</i> | I-1 |